

Table S1.

| Treatment schedule | | Oral Semaglutide | OW Semaglutide | <i>p</i> value |
|-------------------------------------|---|------------------|----------------|----------------|
| Monotherapy | | 1 (0,9%) | 1 (0,6%) | 0,75 |
| Dual Therapy | | 24 (20,9%) | 67 (37,9%) | 0,0022 |
| | Add-on to metformin | 22 (19,15%) | 65 (36,7%) | 0,0016 |
| | Add-on to basal insulin | 0 | 2 (1,2%) | 0,25 |
| | Add-on to SGLT2-i | 2 (1,75%) | 0 | 0,07 |
| Triple Therapy | | 14 (12,2%) | 33 (18,6%) | 0,14 |
| | Add-on Met+Basal | 2 (1,75%) | 16 (9%) | 0,011 |
| | Add-on Met+SGLT2i | 8 (6,95%) | 4 (2,2%) | 0,048 |
| | Add-on Met+Pio | 2 (1,75%) | 6 (3,4%) | 0,4 |
| | Add-on Met+SUs | 2 (1,75%) | 3 (1,7%) | 0,9 |
| | Add-on BB | 0 | 1 (0,6%) | 0,4 |
| | Add-on Met/Empa+Basal | 0 | 3 (1,7%) | 0,16 |
| Replacement therapy | | 76 (66%) | 76 (42,9%) | 0,0001 |
| | DPP4i (\pm SGLT2i o SUs) | 31 (26,95%) | 21 (11,8%) | 0,0016 |
| | SUs | 19 (16,5%) | 11 (6,2%) | 0,0057 |
| | GLP1-RA | 9 (7,8%) | 12 (6,7%) | 0,7 |
| | SGLT2i | 8 (6,95%) | 11 (6,2%) | 0,8 |
| | Metformin | 0 | 2 (1,2%) | 0,25 |
| | Pioglitazone | 4 (3,5%) | 0 | 0,012 |
| | Acarbose | 0 | 1 (0,6%) | 0,4 |
| | Insulin (basal and/or rapid analogues) | 5 (4,3%) | 18 (10,2%) | 0,07 |
| Post-replacement therapy (*) | +Metformin | 33 (43,5%) | 35 (46,1%) | 0,7 |
| | +Basal Insulin | 7 (9,2%) | 5 (6,6%) | 0,5 |
| | +Pioglitazone | 3 (3,9%) | 0 | 0,08 |
| | +SGLT2i | 2 (2,6%) | 0 | 0,15 |
| | +Met+Basal | 11 (14,5%) | 14 (18,5%) | 0,5 |
| | +Met+SGLT2i | 13 (17,1%) | 1 (1,3%) | 0,0008 |
| | +Met+Pio | 1 (1,3%) | 7 (9,2%) | 0,029 |
| | +Met+SUs | 0 | 3 (3,9%) | 0,08 |
| | +SGLT2i+Basal | 0 | 3 (3,9%) | 0,08 |
| | +Met+Pio+SGLT2i | 0 | 1 (1,3%) | 0,3 |
| | Monotherapy | 6 (7,9%) | 5 (6,6%) | 0,7 |
| | +Met+SGLT2i+Basal | 0 | 1 (1,3%) | 0,3 |
| | + BB (\pm Met) | 0 | 1 (1,3%) | 0,3 |

(*)Percentage calculated on the total number of patient who received od/ow semaglutide as replacement therapy