

Burnout among physicians and nursing staff working in the emergency hospital of Tanta University, Egypt

S.A.M. Abdo,¹ R.M. El-Sallamy,¹ A.A.M. El-Sherbiny¹ and I.A. Kabbash¹

الإرهاق لدى الأطباء وطاقم التمريض العاملين في مستشفى الطوارئ بجامعة طنطا، مصر
سناء عبد الفتاح عبده، رانيا مصطفى السلامي، علي علي الشربيني، إبراهيم علي كباش

الخلاصة: لا يُعرف سوى القليل عن الإرهاق المهني لدى العاملين في مجال الرعاية الصحية في مصر. لقد هدفت الدراسة الحالية إلى الكشف عن مدى الإرهاق لدى الأطباء وطاقم التمريض العاملين في مستشفى الطوارئ بجامعة طنطا، وإلى التعرف على بعض العوامل المحددة له. فتم إجراء دراسة مستعرضة شملت جميع الأطباء (ع = 266) وعينة عشوائية منهجية من الممرضات (ع = 284). وتم تقييم الإرهاق باستخدام قائمة ماسلاخ Maslach لتقييم الإرهاق والسلام الفرعية لها. فوجد أن مستوى الإرهاق كان متوسطاً لدى ثلثي المشاركين (66%) ومرتفعاً لدى 24.9% منهم. كما أظهر التحليل متعدد المتغيرات للعوامل التي تؤثر على الإرهاق أن السن والجنس وتواتر التعرض للعنف المتعلق بالعمل وعدد سنوات الخبرة وعبء العمل والإشراف وأنشطة العمل كانت محددات هامة عن الإرهاق لدى المستطلعين. يوصي الباحثون بتدخلات في مجال التثقيف الصحي من خلال برامج تدريب ما قبل التوظيف للوقاية من متلازمة الإرهاق، وإجراء فحص دوري للكشف المبكر عن الإرهاق والقيام بعلاجه.

ABSTRACT Little is known about professional burnout among health-care workers in Egypt. The current study aimed to reveal the extent of burnout among physicians and nursing staff working in the emergency hospital of Tanta University and to identify some of its determinants. A cross-sectional study was carried out on all physicians ($n = 266$) and a systematic random sample of nurses ($n = 284$). Burnout was assessed using the Maslach Burnout Inventory and its subscales. Most of the participants (66.0%) had a moderate level of burnout and 24.9% of them had high burnout. Multivariate analysis of variables affecting burnout showed that age, sex, frequency of exposure to work-related violence, years of experience, work burden, supervision and work activities were significant predictors of burnout among the respondents. The authors recommend health education interventions during pre-employment training programmes for prevention of burnout syndrome and periodic screening for early detection and management of burnout.

L'épuisement professionnel chez des médecins et des personnels infirmiers au service des urgences de l'hôpital universitaire de Tanta (Égypte)

RÉSUMÉ Le syndrome d'épuisement professionnel chez les agents de santé en Égypte est assez mal connu. La présente étude visait à mettre en évidence la fréquence et l'étendue des cas d'épuisement professionnel chez des médecins et personnels infirmiers exerçant aux urgences de l'hôpital universitaire de Tanta et à identifier certains de ses déterminants. Une étude transversale a été menée auprès de tous les médecins ($n = 266$) et d'un échantillon aléatoire systématique des personnels infirmiers ($n = 284$). L'épuisement a été évalué à l'aide du Test d'inventaire de Burnout de Maslach et de ses sous-échelles. La plupart des participants (66,0 %) présentaient un niveau d'épuisement modéré tandis que pour 24,9 % d'entre eux il était élevé. Une analyse multivariée des variables ayant un impact sur l'épuisement professionnel a démontré que l'âge, le sexe, la fréquence des expositions à la violence liée au travail, les années d'expérience, la charge de travail, la supervision et les activités professionnelles étaient des facteurs prédictifs importants de l'épuisement professionnel chez les répondants. Les auteurs recommandent des interventions en matière d'éducation sanitaire au cours de programmes de formation avant l'entrée dans la vie active pour prévenir le syndrome d'épuisement professionnel, puis un dépistage systématique visant la détection précoce et la prise en charge de ce syndrome.

¹Department of Public Health and Community Medicine, Faculty of Medicine, University of Tanta, Tanta, Egypt (Correspondence to I.A. Kabbash: iafkabbash@gmail.com).

Received: 25/12/14; accepted: 28/09/15

Introduction

Burnout is considered an epidemic of modern society and the issue of occupational stress and burnout is receiving increasing attention worldwide. Burnout focuses on specific stressors in the workplace and the environmental pressures affecting the health of employed people (1). It was first described in the mid-1970s by Freudenberger and ever since it has been the subject of many studies (2). Burnout is a psychological term for the experience of long-term exhaustion and diminished interest. Despite this, burnout is not a recognized disorder in the *Diagnostic and Statistical Manual of Mental Disorders* (3).

Burnout syndrome can have a detrimental effect on employee satisfaction, work productivity, mental and physical health, rates of absenteeism and staff turnover and can affect family roles and functions (4). Burnout is an increasing problem among medical staff and is highly prevalent in health-care settings. It is associated with difficult working conditions and feelings of dissatisfaction with work (5). A study among German physicians found that more than one-third of the health-care professionals examined were experiencing professional burnout (6). Another study revealed that burnout can affect up to 45% of medical and nursing staff (7). The impact of burnout not only affects the suffering physicians but extends to their patients if the quality of medical care delivered is undermined (8).

Emergency services are the busiest and most stressful units of hospitals. Their staff might be at risk of burnout syndrome due to inadequate physical working conditions and emotional problems. Staff are under sustained stress owing to the crowded working environment, severity of cases treated and rotational work schedules that disrupt social and family relationships. Furthermore, conflicts with patients' companions, unsafe working environments, work difficulties and critical care

decisions may play a role in the development of burnout syndrome (9).

Little is known about professional burnout among health-care workers in Egypt. The current study was conducted to reveal the prevalence of burnout among physicians and nursing staff working in the emergency hospital of Tanta University and to identify some of the determinants of burnout.

Methods

Study design and setting

A cross-sectional study was conducted over the period 1 November 2012 to 30 April 2013 in the emergency hospital of University of Tanta. This is a 450-bed hospital with the following departments: internal medicine, general surgery, neurosurgery, orthopaedics, burn care unit, toxicology, urology, radiology, intensive care unit and recovery unit. The hospital receives mainly injuries and road traffic accidents from Gharbia governorate and other nearby governorates, in addition to different surgical and medical emergency conditions. The hospital is staffed by 311 physicians who work in different departments in rotating shifts, 490 nurses, 78 head nurses and 70 ancillary workers and technicians. The study was approved by the internal review board of Tanta Faculty of Medicine.

Sampling

The inclusion criteria were physicians and nurses who were in post during the period of the study and consented to participate. The exclusion criteria were any physicians and nurses who had been employed at the hospital for less than 6 months and those who were on vacation during the study period.

The sample size was calculated using *Epi-Info*, version 2002. Based on a level of significance of 95%, 80% power of the study, estimated prevalence of burnout of 50% and error of 5%, the sample size was estimated to be 384. The sample

was increased to 523 to improve the validity of the results.

Out of the total of 311 physicians (146 resident physicians, 19 demonstrators and 146 assistant lecturers) 45 were excluded (42 had less than 6 months in the job and 3 were on vacation). Thus, 266 physicians were eligible to enter the study. Of these, 15 physicians refused to participate and 12 were unreachable. This represents a response rate of 89.8%.

Of the 568 eligible nursing staff, 284 nurses from different departments were included in the study and selected by systematic sampling from the records of employed nurses, after removal of nurses ineligible for the study. The first nurse was chosen randomly from a computerized employment list and then the following every other nurse. The response rate of nurses was 100.0%.

Study tools

The study participants were asked to fill a pre-designed self-administered questionnaire sheet that was designed for this study with sections on sociodemographic, work and health data; job satisfaction; and the assessment of burnout.

Sociodemographic, work and health variables

For the assessment of the determinants of burnout, data were collected about sociodemographic variables (age, sex, residence, marital status, number of family members, number of the rooms in the house, educational level); job characteristics and experiences (job type, department or ward type, years of experience, number of shifts per month, frequency of dealing with and caring for dying patients, previous exposure to work-related violence and its type, and private work outside the hospital); attitudes to work (absenteeism in the last 3 months; thoughts of changing job or moving to a different ward or department in the last 3 months); and health problems (diseases suffered; visits to a

physician or admission to hospital in the last 3 months).

Job satisfaction questionnaire

The job satisfaction questionnaire included 7 main categories as follows:

- General working conditions (4 items about: hours worked each week, flexibility in scheduling, location of work and amount of paid vacation time/sick leave offered).
- Pay and promotion potential (5 items about: salary, opportunities for promotion, benefits, e.g. health insurance, life insurance, job security and recognition for work accomplished).
- Work relationships (3 items about: relationships with co-workers, relationship(s) with supervisor(s) and relationships with subordinates, if applicable).
- Use of skills and abilities (3 items about: opportunity to utilize skills and talents, opportunity to learn new skills and support for additional training and education).
- Work activities (2 items about: variety of job responsibilities, degree of independence associated with work roles and adequate opportunity for periodic changes in duties).
- Work burden (4 items about: workload, time available for family, friends or leisure, work-related stress and administrative burden).
- Supervision (3 items about: method of supervision, whether supervisor admits making errors or not and whether supervisor asks for advice before implementation of tasks).

Respondents were asked to rate how satisfied they were for each item on a 5-point Likert scale from 1 (extremely

dissatisfied) to 5 (extremely satisfied). The satisfaction was arbitrarily classified as: dissatisfied (score < 33.3%); average satisfaction (33.3–66.7%); and satisfied (66.7–100.0%).

Maslach Burnout Inventory

The Maslach Burnout Inventory (MBI), first reported by Maslach et al. in 1996, was the assessment instrument used in the study (10). It has become the almost universally accepted gold standard to assess burnout (11) due to its high reliability and validity (2,10). It was translated into Arabic and then back translated into English, to ensure correct translation of the questions. The Arabic version was revised by 7 experts and tested in a pilot study. The reliability of the questionnaire was tested by Cronbach alpha and found to be 0.81.

The MBI includes 22 items with a 7-point Likert-type rating scale. Items are written in the form of statements about personal feelings or attitudes. The frequency scale is labelled at each point, ranging from 0 (never) to 6 (everyday). The maximum score of the total burnout scale is 132. The MBI has 3 subscales: emotional exhaustion, which refers to both physical and mental exhaustion (statements no. 1–9); depersonalization, which consists of attitude changes of the individual when coming into contact with those who receive his/her services as he/she begins to display a cold and impersonal contact with suffering (statements no. 10–14); and reduced professional accomplishment, which measures perception of the influence of the others, well-being with his/her work, as well as the relationship with problems, evidencing a feeling of dissatisfaction (statements

no. 15–22). Burnout is conceptualized as a continuous variable, ranging from a low to average to high probability of experiencing feelings of each element of burnout. A high degree of burnout is reflected in high scores on the emotional exhaustion (maximum score of 54) and depersonalization subscales (maximum score of 30) and in low scores on the reduced professional accomplishment subscale (maximum score of 48). An average degree of burnout is reflected in average scores on the 3 subscales. A low degree of burnout is reflected in low scores on the emotional exhaustion and depersonalization subscales and in higher scores on the reduced professional accomplishment subscale.

Subscales were classified into low, average and high level of burnout according to Table 1 (12). On the total burnout scale scores of 1–33 are considered as low, 34 – 66 as average and 67–99.9 as high level of burnout.

Data collection

A pilot test on 10% of the subjects (whose results were not included in the study) was carried out before starting data collection to determine potential obstacles, evaluate the adequacy of the designed questionnaire and to estimate the time needed to complete it.

Data were collected by asking the participants to fill the pre-designed self-administered questionnaire. Data collectors visited the hospital 5 days per week and collected an average of 5 questionnaire forms per day. The questionnaire was completed and returned to data collectors in the same session. Written and verbal consent were obtained from subjects prior to their

Table 1 Classification of scores on subscales of the Maslach Burnout Inventory (MBI) (10)

MBI subscales	Level of burnout		
	Low	Average	High
Emotional exhaustion score	< 16	17–26	> 27
Depersonalization score	< 6	7–12	> 13
Reduced professional accomplishment score	> 39	38–32	< 31

participation in the study and those who refused to participate were excluded. Confidentiality and privacy were guaranteed during the whole period of the study by anonymous completion of the questionnaires.

Statistical analysis

Organization, tabulation, presentation and analysis of data were performed using SPSS, version 19. Numerical data were presented as mean and standard deviation (SD). Student t-test, linear correlation and linear regression analysis were used for statistical analysis. Categorical data were presented as numbers and percentages and the chi-squared test was used for statistical analysis. When the chi-squared test was not appropriate, the Monte Carlo exact test was applied. The level of significance adopted was $P < 0.05$.

Results

Demographic data

The questionnaires were completed by 523 staff: 284 nurses and 239 physicians. About one-third of staff had graduated from the secondary school of nursing (31.2%), while 21.6% and 24.3% had graduated from the faculty of medicine with a bachelor of medicine or postgraduate degree respectively (Table 2).

The age range of participants was 20–53 years with a mean of 31.0 (SD 6.0) years. The majority of participants were aged below 40 years (91.8%), female (72.3%) and currently married (73.4%) (Table 1). More than half were from urban areas (57.9%). A total of 42.2% had a family size of 5+. The majority of staff (44.9%) had a crowding index at home of 1–1.5.

Work characteristics

Table 3 shows that 31.0% of physicians dealt with critically ill patients a few times a year, 20.1% once a month and 21.3% once a week. In contrast, 22.5%

Table 2 Sociodemographic characteristics of the studied health-care professionals (n = 523)

Sociodemographic characteristic	No. of respondents	%
Age (years)		
20–	232	44.4
30–	248	47.4
40–	39	7.5
50–	4	0.8
Mean (SD)	31.0 (6.0)	
Sex		
Males	145	27.7
Females	378	72.3
Residence		
Urban	303	57.9
Rural	220	42.1
Marital status		
Single	130	24.9
Married	384	73.4
Divorced	3	0.6
Widow	6	1.1
Family size (no.)		
2	58	11.1
3	114	21.8
4	130	24.9
5	158	30.1
6+	63	12.1
Mean (SD)	4.15 (1.30)	
Crowding index (no. of people/room)		
< 1	105	20.1
1–	235	44.9
1.5–	109	20.8
2–	46	8.8
2.5+	28	5.4
Educational level		
Secondary school (nurse)	163	31.2
High institute (nurse)	69	13.2
Graduate (nurse)	51	9.8
Graduate (physician)	113	21.6
Postgraduate (nurse or physician)	127	24.3
Years of work experience		
< 10	321	61.4
10–	149	28.5
20+	53	10.1
Occupation		
Nurse	233	44.6
Head nurse	51	9.8
Resident	112	21.4
Demonstrator	16	3.1
Assistant lecturer	111	21.2

SD = standard deviation.

Table 3 Distribution of health-care professionals in relation to dealing with critical care patients (near death) and exposure to work-related violence

Variable	Nurses (n = 284)		Physicians (n = 239)		χ^2 -value	P-value
	No.	%	No.	%		
Frequency of dealing with critical patients						
Never	25	8.8	13	5.4	25.93	0.001
Few times a year	46	16.2	74	31.0		
Once monthly	49	17.3	48	20.1		
Once weekly	64	22.5	51	21.3		
Once daily	23	8.1	20	8.4		
More than once daily	77	27.1	33	13.8		
Exposure to work-related violence						
Physical	58	20.4	25	10.5	9.647	0.002
Verbal	272	95.8	219	91.6	3.878	0.049
Sexual	2	0.7	1	0.4		1.000 ^a

^aMonte Carlo exact test.

of nurses dealt with critical patients once a week and 27.1% more than once daily. The difference between nurses and physicians was statistically significant ($P = 0.001$). The majority of physicians and nurses reported that they had been exposed to verbal violence at work (91.6% and 95.8% respectively) ($P = 0.049$). Nurses were

more exposed to physical violence than physicians (20.4% and 10.5% respectively), a difference that was statistically significant ($P = 0.002$). Few respondents reported being exposed to sexual violence at work and the difference between nurses and physicians was not statistically significant ($P = 1.000$).

Levels of burnout

Table 4 shows that 46.9% of respondents scored high on emotional exhaustion, while 44.9% had average levels of depersonalization and a majority (97.7%) had high levels of reduced personal accomplishment. Regarding the total burnout scale, most of the study subjects (66.0%) were classified as

Table 4 Distribution of health-care professionals in relation to scores for total burnout and the 3 subscales of the Maslach Burnout Inventory

Subscale	Level of burnout						χ^2 -value	P-value
	Low		Average		High			
	No.	%	No.	%	No.	%		
Emotional exhaustion							14.28	0.001
Nurses	33	11.6	101	35.6	150	52.8		
Physicians	54	22.6	90	37.7	95	39.7		
Total	87	16.6	191	36.5	245	46.9		
Depersonalization							29.57	0.001
Nurses	137	48.2	126	44.4	21	7.4		
Physicians	76	31.8	109	45.6	54	22.6		
Total	213	40.7	235	44.9	75	14.4		
Reduced personal accomplishment							4.171	0.041
Nurses	0	0.0	10	3.5	274	96.5		
Physicians	0	0.0	2	0.8	237	99.2		
Total	0	0.0	12	2.3	511	97.7		
Total burnout scale							2.315	0.314
Nurses	22	7.7	186	65.5	76	26.8		
Physicians	26	10.9	159	66.5	54	22.6		
Total	48	9.1	345	66.0	130	24.9		

having a moderate level of burnout and 24.9% as having high burnout.

More than half of nurses (52.8%) experienced high emotional exhaustion compared with 39.7% of physicians, a difference that was statistically significant ($P < 0.001$). For the depersonalization subscale, 48.2% of nurses had low scores and 44.4% moderate scores, compared with 31.8% and 45.6% of physicians respectively, and the difference was statistically significant ($P < 0.001$). Almost all the physicians (99.2%) experienced highly reduced feelings of personal accomplishment and the rate was slightly lower among nurses (96.5%) ($P = 0.041$). Regarding the total burnout scale, the difference between nurses and physicians was not statistically significant.

Job satisfaction

Table 5 shows that the majority of nurses and physicians had average levels of job satisfaction (89.4% and 81.6% respectively), a difference that was statistically significant ($P = 0.028$). For general work conditions, 83.8% of nurses experienced average satisfaction compared with 61.9% of physicians ($P = 0.001$). Concerning promotion and financial aspects, 71.1% of nurses experienced average satisfaction compared with 74.9% of physicians ($P = 0.001$). For relationships in work, 64.8% of nurses were satisfied with their relationships with their colleagues, supervisors and subordinates compared with 60.3% of physicians ($P = 0.019$). Fewer nurses had an average satisfaction level regarding using skills and abilities (51.4%)

compared with physicians (69.9%) ($P = 0.020$). Regarding work activities, the majority of nurses and physicians (62.7% and 68.6% respectively) experienced average satisfaction and around one-fifth were dissatisfied (21.8% and 21.8% respectively) ($P = 0.020$). The majority of nurses (69.0%) and more than half of physicians (56.9%) had average satisfaction regarding work burden. On the other hand, 28.5% and 38.5% of nurses and physicians respectively were dissatisfied with the work burden ($P = 0.013$). As regards supervision at work, 45.8% of nurses had average satisfaction compared with 56.9% of physicians ($P = 0.001$).

Predictors of burnout

Multivariate analysis by linear regression revealed that age, frequency of

Table 5 Distribution of health-care professionals in relation to level of job satisfaction

Item	Level of job satisfaction						χ^2 -value	P-value
	Dissatisfied		Average		Satisfied			
	No.	%	No.	%	No.	%		
General work condition								
Nurse	17	6.0	238	83.8	29	10.2	38.21	0.001
Physician	55	23.0	148	61.9	36	15.1		
Promotion and financials								
Nurse	64	22.5	202	71.1	18	6.3	21.86	0.001
Physician	24	10.0	179	74.9	36	15.1		
Relationships in work								
Nurse	4	1.4	96	33.8	184	64.8	7.891	0.019
Physician	14	5.9	81	33.9	144	60.3		
Using skills and abilities								
Nurse	72	25.4	146	51.4	66	23.2	19.80	0.001
Physician	44	18.4	167	69.9	28	11.7		
Work activities								
Nurse	62	21.8	178	62.7	44	15.5	7.873	0.020
Physician	57	23.8	164	68.6	18	7.5		
Work burden								
Nurse	81	28.5	196	69.0	7	2.5	8.624	0.013
Physician	92	38.5	136	56.9	11	4.6		
Supervision at work								
Nurse	54	19.0	130	45.8	100	35.2	22.84	0.001
Physician	63	26.4	136	56.9	40	16.7		
Total job satisfaction								
Nurse	11	3.9	245	89.4	19	6.7	7.123	0.028
Physician	20	8.4	195	81.6	24	10.0		

work-related exposure to violence, years of experience, work burden, supervision and work activities were the significant predictors for burnout (Table 6).

Discussion

Occupational-related burnout is increasingly recognized as a serious problem affecting many people working in the human services, particularly health-care workers, who suffer from negative feelings for self, work and life (13). Health-care professionals work with others in emotionally demanding situations and are exposed to their clients' psychological, socio-economic and physical problems. As a result, burnout can easily develop, especially in sectors where critical care is provided (14). Burnout has been associated with absenteeism from work, ineffectiveness, interpersonal conflicts, lower productivity, job dissatisfaction, reduced organizational commitment and staff turnover. It predicts increased rates of illness, fatigue, substance misuse, depression, anxiety and irritability (15).

The aim of this study was to assess the prevalence of burnout syndrome among health-care workers of the emergency hospital, University of Tanta and to identify some of its determinants. About two-thirds of our study subjects had a moderate level of burnout and about one-quarter had a high level of burnout. This may be due to the overburdened health-care system in Egypt, especially the emergency sector, understaffing, especially among nursing staff, lack of resources, inadequate salaries, lack of control, difficult work schedules with long hours of work and frequent shifts, inadequate security and poor career advancement.

The results of studies on burnout are controversial, with some studies revealing alarmingly high rates of burnout among physicians, while other studies finding much lower rates. In

Table 6 Multivariate analysis of variables affecting scores for total burnout on the Maslach Burnout Inventory among the studied health-care professionals

Variable	t-value	P-value
Age	5.097	0.001
Frequency of exposure to work-related violence	4.567	0.001
Years of experience	3.969	0.001
Work burden	3.960	0.001
Supervision	2.669	0.008
Work activities	2.570	0.010

Egypt, Yousef et al. in their study in Suez Canal University about the prevalence of burnout syndrome among residents of the university hospitals showed that 63.1% of the responding residents met the criteria for burnout (12). Elsewhere in the Eastern Mediterranean Region, Halayem-Dhouib et al. in Tunisia found that burnout syndrome was highly prevalent among nurses and medical residents (16). In Saudi Arabia, Al-Turki et al. reported that the prevalence of burnout syndrome among multinational nurses was high (17). In contrast, a study carried out in Qatar found that 12.6% of the responding primary health-care physicians were experiencing burnout (18).

Previous European studies using the MBI reported widely varying burnout rates among medical and paramedical teams, ranging from 15–82% (19,20). The prevalence of burnout among staff of the departments of anaesthesia and critical care medicine was about 30% of nurses and 40–50% of physicians (21). Cydulka and Korte mentioned that previous estimates of burnout among emergency physicians in the United States of America range from 25–60% and more than 90% among Canadian emergency physicians (22).

In African countries, high rates of burnout among maternal health staff at a referral hospital in Malawi have been reported (23). Other studies performed in Nigeria among health professionals (24) and in a Kenyan psychiatric hospital (25) have described high rates of burnout.

In South America, most studies on burnout syndrome were from Brazil. The prevalence of burnout syndrome among community-based health agents in the city of Sao Paulo was been estimated to be 24% (26). Another Brazilian study performed on nurses in Santa Catarina showed that 35.7% displayed burnout (27).

In the last few years, several Asian studies have focused on burnout syndrome. From Singapore, Lim et al. reported that Singaporean nurses experienced high levels of stress related to work and emergency care (28). In Mongolia, a study by Bagaajav et al. detected an increasing level of stress among medical doctors. Excessive workload was a significant source of developing burnout, job stress and job dissatisfaction (29).

The Australian Medical Association performed a health survey on 914 Australian and New Zealand junior doctors and found that 69% had burnout symptoms (30).

The difference in the prevalence of burnout syndrome across different countries can possibly be explained by variations in culture, the nature of the health system (including structural and service delivery problems in different countries), patients' attitudes and the role of physicians as health-care providers. There are also differences in the assessment scales and study designs used in the various studies of burnout syndrome (31).

Almost half of our respondents (46.9%) showed high emotional

exhaustion, 97.7% showed highly reduced personal accomplishment and 14.4% showed high depersonalization. The subscale of reduced personal accomplishment reflects lower competence at work, which in turn affects the effectiveness of work leading to decreased quality of health-care services delivered.

Kalemoglu and Keskin in Istanbul, Turkey, reported that among emergency service personnel, 44.7% showed high emotional exhaustion, 33.2% high depersonalization and 28% high reduced professional accomplishment (9). A French study by Embriaco et al. found a high level of emotional exhaustion in 19% of intensive care unit staff, high depersonalization in 37% and highly reduced professional accomplishment in 39% (7). A Belgian study of health-care workers managing chronic patients with consciousness disorders reported that 33% showed moderate to severe emotional exhaustion, 36% showed moderate to severe depersonalization and only 3% presented low personal accomplishment (32).

Our study revealed that more than a half of nurses experienced high emotional exhaustion compared with physicians. Physicians had higher level of depersonalization than nurses and most of the physicians and nurses showed high reduced personal accomplishment. This is consistent with Gosses et al., who reported that nurses were more emotionally exhausted than physicians. However, the rates of depersonalization and reduced personal accomplishment were greater in nurses even though the difference was not statistically significant (32). On the other hand, Escriba-Aguir et al., in their study on emergency medical and nursing staff in Spain, found that the prevalence of emotional exhaustion and reduced professional accomplishment was higher among doctors than nurses (33). Nursing staff are most likely to develop burnout due to the nature of their job, with many tasks, greater proximity and

time spent with patients and relatives, and the emotional demands of their work (34).

Rajeev and Rashmi Khanna in Rajasthan in India found that 29.2% of medical professional showed high levels of emotional exhaustion, 20.0% showed high depersonalization and 17.9% showed high reduced professional accomplishment (35). Among medical residents, Prins et al. found that rates of emotional exhaustion were 41–50% and depersonalization 34–70% (36), while in another study including nurses from different specialties, Sahraian et al. found reported rates of 23% for emotional exhaustion and 5% for depersonalization. (34). In contrast, Abdulghafour et al. revealed that physicians in primary health-care centres in Kuwait had a relatively low mean percentage scores for emotional exhaustion and depersonalization, but they had high mean scores for personal accomplishment (8).

A great challenge encountered during emergency service is that the care of critical patients, such as undiagnosed cases, trauma patients with crush injuries and critical or dying patients, might lead to psychological depression in emergency service workers (9). Our study demonstrated a weak significant correlation between burnout and dealing with dying patients. Likewise, Poncet et al. (37) and Embriaco et al. (7) revealed that severe burnout was associated with end-of-life factors such as caring for dying patients.

Nurses in our study were more exposed to physical and verbal violence than were physicians. Kalemoglu and Keskin also demonstrated a high level of aggressive behaviour by patients and their companions in the emergency service. These aggressive acts might comprise personal threats or even use of weapons. This might complicate the safety of doctors, nurses and personnel taking care of patients and facilitate the development of burnout (9).

Multivariate analysis of variables affecting burnout in our study revealed that age, frequency of exposure to violence at work, years of experience, work burden, supervision and work activities were significant predictors of burnout syndrome among the study population. In Turkey, Demir et al. reported that problems with childcare were significantly associated with burnout among Turkish nurses (38). Lagerström et al. also reported work dissatisfaction, health threats and disequilibrium between family and work demands as the most important factors contributing to burnout among Iranian nurses (39).

A study on Cypriot physiotherapists in 2010 reported that regression analysis showed that the perception of stressful job, low salary, age group and employment sector predicted high burnout scores (14). Thomas and Valli in South Africa, reported that public hospital physicians had high levels of burnout syndrome and the main sources of stress were understaffing, lack of resources, difficult work schedules, inadequate job security and poor salaries. In addition, senior doctors showed lower job satisfaction (40).

The Australian Medical Association performed a health survey with 914 Australian and New Zealand junior doctors. Approximately 71% had low job satisfaction and 69% had burnout symptoms and half of them reported that their workload was excessive (30). Meanwhile, Lee et al. found that physician burnout was negatively associated with autonomy and positive work attitudes. It was positively associated with workload, constraining organizational structure, incivility, conflicts, violence, low quality and safety standards, negative work attitudes, work-life conflict and contributors to poor mental health (41).

One of the limitations of our study is that a cross-sectional study is not considered the ideal tool to study the causes of burnout syndrome.

Conclusion and recommendations

The present study in the emergency hospital of University of Tanta, Egypt concluded that about one-quarter of study nurses and physicians (26.8% and 22.6%) suffered from high levels of

burnout syndrome. Age, frequency of exposure to violence at work, years of experience, work burden, supervision and work activities were significant predictors of burnout syndrome among the health-care professionals studied. We recommend a longitudinal study to establish the causation of burnout syndrome;

health education interventions during pre-employment training programmes for prevention of burnout syndrome; and periodic screening during work for early detection and management of burnout among health-care professionals.

Finding: None.

Competing interests: None declared.

References

- Schaufeli WB, Greenglass ER. Introduction to special issue on burnout and health. *Psychol Health*. 2001 Sep;16(5):501-10. PMID:22804495
- Maslach C, Schaufeli WB, Leiter MP. Job burnout. *Annu Rev Psychol*. 2001;52:397-422. PMID:11148311
- Kraft U. Burned out. *Sci Am Mind*. 2006; (June/July):28-33.
- Tennant C. Work-related stress and depressive disorders. *J Psychosom Res*. 2001 Nov;51(5):697-704. PMID:11728512
- De Oliveira GS Jr, Ahmad S, Stock MC, Harter RL, Almeida MD, Fitzgerald PC, et al. High incidence of burnout in academic chairpersons of anesthesiology: should we be taking better care of our leaders? *Anesthesiology*. 2011 Jan;114(1):181-93. PMID:21178674
- Janus K, Amelung VE, Gaitanides M, Schwartz FW. German physicians "on strike"—shedding light on the roots of physician dissatisfaction. *Health Policy*. 2007 Aug;82(3):357-65. PMID:17137674
- Embriaco N, Azoulay E, Barrau K, Kentish N, Pochard F, Loundou A, et al. High level of burnout in intensivists: prevalence and associated factors. *Am J Respir Crit Care Med*. 2007 Apr 1;175(7):686-92. PMID:17234905
- Abdulghafour YA, Bo-hamrab AM, Al-Randic MS, Kamel MI, El-Shazly MK. Burnout syndrome among physicians working in primary health care centers in Kuwait. *Alexandria Journal of Medicine*. 2011;47:351-7.
- Kalemoglu M, Keskin O. Burnout syndrome at the emergency service. *Scand J Trauma Resusc Emerg Med*. 2006;14:37-40.
- Maslach C, Jackson SE, Leiter MP. *Maslach Burnout Inventory manual*. 3rd ed. Palo Alto (CA): Consulting Psychologist Press; 1996.
- Schuttler N, Toppinen S, Kalimo R, Schaufeli W. The factorial validity of the Maslach Burnout Inventory across occupational groups and nations. *J Occup Organ Psychol*. 2000;73:53-66.
- Yousef IM, Hosny AO, Elsayed OI, Ali EG. Burnout Syndrome among Resident Physician in Suez Canal University Hospital. *Current Psychiatry Ain Shams University*. 2006 March;13(1):24-43.
- Sørgaard KW, Ryan P, Hill R, Dawson I; OSCAR Group. Sources of stress and burnout in acute psychiatric care: inpatient vs. community staff. *Soc Psychiatry Psychiatr Epidemiol*. 2007 Oct;42(10):794-802. PMID:17700977
- Pavakis A, Raftopoulos V, Theodorou M. Burnout syndrome in Cypriot physiotherapists: a national survey. *BMC Health Serv Res*. 2010;10:63. PMID:20222948
- Knudsen HK, Ducharme LJ, Roman PM. Clinical supervision, emotional exhaustion, and turnover intention: a study of substance abuse treatment counselors in the Clinical Trials Network of the National Institute on Drug Abuse. *J Subst Abuse Treat*. 2008 Dec;35(4):387-95. PMID:18424048
- Halayem-Dhouib S, Zaghoudi L, Zremdini R, Maalej I, Béchir MB, Labbène R. Burnout en psychiatrie : une expérience tunisienne [Burnout among mental health professionals: a Tunisian experience]. *Rev Epidemiol Sante Publique*. 2010 Dec;58(6):403-8. PMID:21094003
- Al-Turki HA, Al-Turki RA, Al-Dardas HA, Al-Gazal MR, Al-Maghrabi GH, Al-Enizi NH, et al. Burnout syndrome among multinational nurses working in Saudi Arabia. *Ann Afr Med*. 2010 Oct-Dec;9(4):226-9. PMID:20935422
- Abdulla L, Al-Qahtani DM, Al-Kuwari MG. Prevalence and determinants of burnout syndrome among primary healthcare physicians in Qatar. *South African Family Practice Journal*. 2011;53(4):380-3.
- Hyman SA, Michaels DR, Berry JM, Schildcrout JS, Mercaldo ND, Weinger MB. Risk of burnout in perioperative clinicians: a survey study and literature review. *Anesthesiology*. 2011 Jan;114(1):194-204. PMID:21178675
- Roth M, Morrone K, Moody K, Kim M, Wang D, Moadel A, et al. Career burnout among pediatric oncologists. *Pediatr Blood Cancer*. 2011 Dec 15;57(7):1168-73. PMID:21548010
- Wallace JE, Lemaire JB, Ghali WA. Physician wellness: a mishap quality indicator. *Lancet*. 2009 Nov 14;374(9702):1714-21. PMID:19914516
- Cydulka RK, Korte R. Career satisfaction in emergency medicine: the ABEM Longitudinal Study of Emergency Physicians. *Ann Emerg Med*. 2008 Jun;51(6):714-722.e1. PMID:18395936
- Thorsen VC, Tharp AL, Meguid T. High rates of burnout among maternal health staff at a referral hospital in Malawi: A cross-sectional study. *BMC Nurs*. 2011;10:9. PMID:21605379
- Onyilezugbo E, Nawfor C. Construct validation of nurse's self-concept questionnaire in Nigeria. *Eur J Soil Sci*. 2010;15:467-74.
- Ndetei DM, Pizzo M, Maru H, Ongecha FA, Khasakhala LI, Mutiso V, et al. Burnout in staff working at the Mathari psychiatric hospital. *Afr J Psychiatry (Johannesbg)*. 2008 Aug;11(3):199-203. PMID:19588043
- Silva AT, Menezes PR. Burnout syndrome and common mental disorders among community-based health agents. *Rev Saude Publica*. 2008 Oct;42(5):921-9. PMID:18833390
- Moreira DdeS, Magnago RF, Sakae TM, Magajewski FR. Prevalência da síndrome de burnout em trabalhadores de enfermagem de um hospital de grande porte da Região Sul do Brasil. [Prevalence of burnout syndrome in nursing staff in a large hospital in south of Brazil]. *Cad Saude Publica*. 2009 Jul;25(7):1559-68. PMID:19578577
- Lim J, Bogossian F, Ahern K. Stress and coping in Singaporean nurses: a literature review. *Nurs Health Sci*. 2010 Jun;12(2):251-8. PMID:20602699
- Bagaajav A, Myagmarjav S, Nanjid K, Otgon S, Chae YM. Burnout and job stress among mongolian doctors and nurses. *Ind Health*. 2011;49(5):582-8. PMID:21804269

30. Markwell AL, Wainer Z. The health and wellbeing of junior doctors: insights from a national survey. *Med J Aust.* 2009 Oct 19;191(8):441-4. PMID:19835538
31. Kolstad HA, Hansen AM, Kærgaard A, Thomsen JF, Kaerlev L, Mikkelsen S, et al. Job strain and the risk of depression: is reporting biased? *Am J Epidemiol.* 2011 Jan 1;173(1):94-102. PMID:21071605
32. Gosseries O, Demertzi A, Ledoux D, Bruno MA, Vanhaudenhuyse A, Thibaut A, et al. Burnout in healthcare workers managing chronic patients with disorders of consciousness. *Brain Inj.* 2012;26(12):1493-9. PMID:22725684
33. Escribà-Agüir V, Martín-Baena D, Pérez-Hoyos S. Psychosocial work environment and burnout among emergency medical and nursing staff. *Int Arch Occup Environ Health.* 2006 Nov;80(2):127-33. PMID:16710712
34. Sahraian A, Fazelzadeh A, Mehdizadeh AR, Toobae SH. Burnout in hospital nurses: a comparison of internal, surgery, psychiatry and burns wards. *Int Nurs Rev.* 2008 Mar;55(1):62-7. PMID:18275537
35. Khanna R, Khanna R. Is medicine turning into unhappy profession? *Indian J Occup Environ Med.* 2013 Jan;17(1):2-6. PMID:24082640
36. Prins JT, Gazendam-Donofrio SM, Tubben BJ, van der Heijden FM, van de Wiel HB, Hoekstra-Weebers JE. Burnout in medical residents: a review. *Med Educ.* 2007 Aug;41(8):788-800. PMID:17661887
37. Poncet MC, Toullic P, Papazian L, Kentish-Barnes N, Timsit JF, Pochard F, et al. Burnout syndrome in critical care nursing staff. *Am J Respir Crit Care Med.* 2007 Apr 1;175(7):698-704. PMID:17110646
38. Demir A, Ulusoy M, Ulusoy MF. Investigation of factors influencing burnout levels in the professional and private lives of nurses. *Int J Nurs Stud.* 2003 Nov;40(8):807-27. PMID:14568363
39. Lagerström M, Josephson M, Arsalani N, Fallahi-Khoshknab M. Striving for balance between family and work demands among Iranian nurses. *Nurs Sci Q.* 2010 Apr;23(2):166-72. PMID:20378918
40. Thomas LS, Valli A. Levels of occupational stress in doctors working in a South African public-sector hospital. *S Afr Med J.* 2006 Nov;96(11):1162-8. PMID:17167700
41. Lee RT, Seo B, Hladkyj S, Lovell BL, Schwartzmann L. Correlates of physician burnout across regions and specialties: a meta-analysis. *Hum Resour Health.* 2013;11(1):48. PMID:24074053