

Online Resource 2

Article title: Structural damage progression in patients with early rheumatoid arthritis treated with methotrexate, baricitinib or baricitinib plus methotrexate based on clinical response in the phase 3 RA-BEGIN study

Journal: *Clinical Rheumatology*

Authors: Désirée van der Heijde, Patrick Durez, Georg Schett, Esperanza Naredo, Mikkel Østergaard, Gabriella Meszaros, Francesco De Leonardi, Inmaculada de la Torre, Pedro López-Romero, Douglas Schlichting, Eric Nantz, Roy Fleischmann

Affiliation of corresponding author: Leiden University Medical Center, Leiden, The Netherlands, email address: mail@dvanderheijde.nl

Patient baseline characteristics grouped by SDAI score and treatment (SDAI-Group A: sustained SDAI score ≤ 11 at weeks 16, 20 and 24; SDAI-Group B: SDAI score > 11 or missing data at any of weeks 16, 20 and 24)

Treatment (N; % of treatment group)	SDAI-Group A (N=209)			SDAI-Group B (N=375)		
	MTX (N=53; 25.2%)	Bari (N=62; 39.0%)	Bari + MTX (N=94; 43.7%)	MTX (N=157; 74.8%)	Bari (N=97; 61.0%)	Bari + MTX (N=121; 56.3%)
	52 \pm 13	51 \pm 13	45 \pm 14	50 \pm 14	51 \pm 13	51 \pm 13
Age, years	52 \pm 13	51 \pm 13	45 \pm 14	50 \pm 14	51 \pm 13	51 \pm 13
Female	36 (67.9)	47 (75.8)	67 (71.3)	112 (71.3)	74 (76.3)	89 (73.6)
BMI, kg/m ²	26.3 \pm 6.0	27.7 \pm 7.4	25.7 \pm 5.1	26.6 \pm 6.3	26.3 \pm 6.1	27.3 \pm 6.5
Smoker	10 (18.9)	17 (27.4)	20 (21.3)	38 (24.2)	20 (20.6)	27 (22.3)
Duration of RA from diagnosis, years ^a	0.2 (0.1, 0.4)	0.2 (0.1, 0.4)	0.2 (0.1, 1.0)	0.2 (0.1, 0.7)	0.4 (0.1, 1.6)	0.2 (0.1, 1.1)
ACPA positive ^b	48 (90.6)	59 (95.2)	88 (93.6)	145 (92.9)	83 (85.6)	104 (86.0)
RF positive ^c	50 (94.3)	61 (98.4)	88 (93.6)	153 (97.5)	94 (96.9)	116 (95.9)
≥ 1 joint erosion	35 (66.0)	41 (66.1)	52 (55.3)	103 (66.0)	64 (66.0)	85 (70.2)
hsCRP, mg/L	16.4 \pm 12.7	19.4 \pm 19.2	25.7 \pm 28.4	24.4 \pm 23.8	26.5 \pm 29.6	23.2 \pm 30.3
DAS28-hsCRP	5.7 \pm 0.9	5.6 \pm 0.9	5.8 \pm 0.9	5.9 \pm 1.0	6.1 \pm 1.0	6.0 \pm 0.9
SDAI	39.7 \pm 12.9	38.2 \pm 12.4	40.9 \pm 12.8	42.3 \pm 14.2	45.6 \pm 14.2	44.2 \pm 13.6
mTSS	9.7 \pm 21.8	10.5 \pm 25.5	7.5 \pm 14.7	12.6 \pm 22.4	15.2 \pm 28.0	14.9 \pm 23.5
HAQ-DI total score	1.5 \pm 0.7	1.5 \pm 0.8	1.6 \pm 0.7	1.7 \pm 0.7	1.7 \pm 0.7	1.6 \pm 0.6

Data are reported as mean \pm standard deviation or n (%) unless otherwise indicated

^aData presented as medians with interquartile range

^bACPA positive (>ULN [ULN=10 U/mL])

^cRF positive (>ULN [ULN=14 U/mL])

ACPA anti-cyclic citrullinated peptide antibody, *Bari* baricitinib, *BMI* body mass index, *DAS28* Disease Activity Score for 28-joint counts, *HAQ-DI* Health Assessment Questionnaire-Disability Index, *hsCRP* high-sensitivity C-reactive protein, *mTSS* van der Heijde-modified total Sharp score, *MTX* methotrexate, *RA* rheumatoid arthritis, *RF* rheumatoid factor, *SDAI* Simplified Disease Activity Index, *ULN* upper limit of normal