SUPPLEMENTAL MATERIAL

Phase 1 Study of Pembrolizumab Plus Chemotherapy in Japanese Patients With Extensive-Stage Small-Cell Lung Cancer

Naoyuki Nogami,¹ Takaaki Tokito,² Yoshitaka Zenke,³ Miyako Satouchi,⁴ Takashi Seto,⁵ Hideo Saka,⁶ Junko Otani,⁷ Shirong Han,⁷ Kazuo Noguchi,⁷ Makoto Nishio⁸

¹Ehime University Graduate School of Medicine, Department of Community Medicine, Pulmonology and Cardiology, Toon, Japan; ²Division of Respirology, Neurology, and Rheumatology, Department of Internal Medicine, Kurume University School of Medicine, Kurume, Japan; ³Department of Thoracic Oncology, National Cancer Center Hospital East, Kashiwa, Japan; ⁴Department of Thoracic Oncology, Hyogo Cancer Center, Akashi, Japan; ⁵Department of Thoracic Oncology, National Hospital Organization Kyushu Cancer Center, Fukuoka, Japan; ⁶Department of Respiratory Medicine, National Hospital Organization Nagoya Medical Center, Nagoya, Japan; ⁷MSD K.K., Tokyo, Japan; ⁸Department of Thoracic Medical Oncology, The Cancer Institute Hospital of Japanese Foundation for Cancer Research, Tokyo, Japan

Corresponding Author:	Naoyuki Nogami
	Department of Community Medicine, Pulmonology and
	Cardiology
	Ehime University Graduate School of Medicine
	454 Shitsukawa, Toon, Ehime 791-0295
	Japan
	Email: nogami.naoyuki.zx@ehime-u.ac.jp

Table S1. Subsequent anticancer therapy

	Second line	Third line	Fourth line
Subsequent therapy	(n = 13)	(n = 8)	(n = 5)
Amrubicin	10	3	0
Cisplatin plus etoposide	1	0	0
Carboplatin plus etoposide	1	0	0
Investigational drug	1	1	0
Topotecan	0	2	0
Cisplatin plus irinotecan	0	2	0
Eribulin	0	0	2
Carboplatin plus paclitaxel	0	0	1
Erlotinib	0	0	1
Atezolizumab, carboplatin, and etoposide	0	0	1