

TABLE 1 Summary of case reports of NPHF infections in XLA (Total 27cases) .

|                   |                                        | Cases                                 | (%) |       | Sample                | Cases                | (%)       |       |       |
|-------------------|----------------------------------------|---------------------------------------|-----|-------|-----------------------|----------------------|-----------|-------|-------|
| Species           | single pathogen                        | H. Cinaedi                            | 11  | 40.74 | 16S sequencing        | Blood                | 2         | 7.41  |       |
|                   |                                        | H. bilis                              | 7   | 25.92 | (14cases, 51.85%)     | Biopsy               | 6         | 22.22 |       |
|                   |                                        | H.canis                               | 1   | 3.70  |                       | Pus                  | 1         | 3.70  |       |
|                   |                                        | H. fennelliae                         | 1   | 3.70  |                       | Blood                | 5         | 18.52 |       |
|                   |                                        | Helicobacter sp                       | 1   | 3.70  | PCR                   | -                    | 3         | 11.11 |       |
|                   | coinfection                            | Helicobacter/Flexispira-like organism | 1   | 3.70  | Diagnosis method      | mNGS (*)             | Blood (*) | 2     | 7.41  |
|                   |                                        | H. Equorum -like                      | 1   | 3.70  |                       | Pus (*)              | 2         | 7.41  |       |
|                   |                                        | Canadensis-like                       | 1   | 3.70  |                       | Cultures             | Blood     | 4     | 14.81 |
|                   |                                        | Helicobacter trogontum                | 1   | 3.70  |                       | Pus                  | 0         | 0.00  |       |
|                   |                                        | H. Canis and H. Bilis (F. rappini)    | 1   | 3.70  | 23S sequencing        | -                    | 1         | 3.70  |       |
| Type of infection | cellulitis (*)<br>(17cases, 62.96%)    | H.equorum and H.canadensis (*)        | 1   | 3.70  | Mass spectrometry     | -                    | 5         | 18.52 |       |
|                   |                                        | Lower limbs (*)                       | 15  | 55.55 | Imipenem/cilastin (*) | -                    | 8         | 29.63 |       |
|                   |                                        | upper limb                            | 4   | 14.81 | Meropenem             | -                    | 5         | 18.52 |       |
|                   | P-G-like ulcer (*)<br>(7cases, 25.92%) | unknown                               | 2   | 7.41  | Antibiotics           | Ertapenem            | -         | 3     | 11.11 |
|                   |                                        | lower limbs (*)                       | 6   | 22.22 |                       | Panipenem/betamipron | -         | 1     | 3.70  |
|                   |                                        | upper limbs                           | 1   | 3.70  |                       | Ceftazidime          | -         | 1     | 3.70  |

TABLE 1 continue

|                   |                                                    |           |    |       |                                                   |                |   |       |       |
|-------------------|----------------------------------------------------|-----------|----|-------|---------------------------------------------------|----------------|---|-------|-------|
|                   | Hyperpigmented plaques<br>and subcutaneous nodules | both legs | 1  | 3.70  | Doxycycline (*)                                   | -              | 7 | 25.93 |       |
|                   | Erythematous plaques                               | ankle     | 1  | 3.70  | Minocycline                                       | -              | 6 | 22.22 |       |
|                   | * skin infection                                   | -         | 19 | 70.37 | Amoxicillin/amoxicillin/clavula<br>nate potassium | -              | 4 | 14.81 |       |
|                   | (all the above)                                    | -         |    |       | Netronidazole                                     | -              | 4 | 14.81 |       |
|                   | Sepsis (*)                                         | -         | 12 | 44.44 | Ciprofloxacin                                     | -              | 1 | 3.70  |       |
|                   | Osteomyelitis                                      | -         | 2  | 7.41  | Tobramycin                                        | -              | 1 | 3.70  |       |
|                   | Suppurative cholangitis                            | -         | 3  | 11.11 | Rifampicin (*)                                    | -              | 3 | 11.11 |       |
| Type of infection | Abdominal abscess                                  | -         | 1  | 3.70  | Antibiotics                                       | Clarithromycin | - | 3     | 11.11 |
|                   | Refractory chronic pleurisy                        | -         | 1  | 3.70  | Levofloxacin                                      | -              | 1 | 3.70  |       |
|                   | synovitis                                          | -         | 1  | 3.70  | Amikacin                                          | -              | 1 | 3.70  |       |
|                   |                                                    |           |    |       | Moxifloxacin                                      | -              | 2 | 7.41  |       |
|                   |                                                    |           |    |       | Gentamicin                                        | -              | 5 | 18.52 |       |
|                   |                                                    |           |    |       | Ceftriaxone                                       | -              | 3 | 11.11 |       |
|                   |                                                    |           |    |       | Cefepime                                          | -              | 1 | 3.70  |       |
|                   |                                                    |           |    |       | TrimethoprimSulfamethoxazole                      | -              | 2 | 7.41  |       |
|                   |                                                    |           |    |       | Vancomycin                                        | -              | 1 | 3.70  |       |
|                   |                                                    |           |    |       | Linezolid                                         | -              | 1 | 3.70  |       |
|                   |                                                    |           |    |       | Kanamycin                                         | -              | 1 | 3.70  |       |
|                   |                                                    |           |    |       | Fosfomycin (*)                                    | -              | 2 | 7.41  |       |
|                   |                                                    |           |    |       | Azithromycin                                      | -              | 2 | 7.41  |       |

(\*) : our patient \* There are 4 patients have both cellulitis and PG-like ulcers, and 4patients have both upper and lower extremity cellulitis.

## References

- [1]Weir S, Cuccherini B, Whitney AM, et al. Recurrent Bacteremia Caused by a “Flexispira”-Like Organism in a Patient With X-Linked (Bruton’s) Agammaglobulinemia. *J. Clin. Microbiol.* 1999;37, 2439 – 2445.
- [2] Cuccherini B, Chua K, Gill V, et al. Bacteremia and Skin/Bone Infections in Two Patients With X-Linked Agammaglobulinemia Caused by an Unusual Organism Related to *Flexispira/Helicobacter* Species. *Clin. Immunol.* 2000;97, 121-129.
- [3]Han S, Schindel C., Genitsariotis R, et al. Identification of a Unique *Helicobacter* Species by 16S rRNA Gene Analysis in an Abdominal Abscess From a Patient With X-Linked Hypogammaglobulinemia. *J. Clin. Microbiol.* 2000;38, 2740 – 2742.
- [4]Gerrard J, Alfredson D, and Smith I. Recurrent Bacteremia and Multifocal Lower Limb Cellulitis Due to *Helicobacter*-Like Organisms in a Patient With X-Linked Hypogammaglobulinemia. *Clin. Infect. Dis.* 2001;33, E116 – E118.
- [5]Simons E, Spacek L A, Lederman HM, and Winkelstein J A. *Helicobacter Cinaedi* Bacteremia Presenting as Macules in an Afebrile Patient With X-Linked Agammaglobulinemia. *Infection*;2004;32, 367 – 368.
- [6]Leemann C, Gambillara E, Prod'hom G ,et al. First case of bacteremia and multifocal cellulitis due to *Helicobacter canis* in an immunocompetent patient. *J. Clin. Microbiol.* 2006 ;44(12):4598-600.
- [7] Schwarze-Zander C, Becker S, Wenzel J,et al. Bacteremia Caused by a Novel *Helicobacter* Species in a 28-Year-Old Man Wi[th X-Linked Agammaglobulinemia. *J. Clin. Microbiol.* 2010;48,4672 – 4676.
- [8]Murray P R, A Uzel, G Ranken, et al. Pyoderma Gangrenosum-Like Ulcer in a Patient With X-Linked Agammaglobulinemia: Identification of *Helicobacter Bilis* by Mass Spectrometry Analysis. *Arch.*2010;146, 523–526. doi: 10.1001/archdermatol.2010.86
- [9]Funato M, Kaneko H, Ohkusu K, et al.. Refractory Chronic Pleurisy Caused by *Helicobacter Equorum*-Like Bacterium in a Patient With X-Linked Agammaglobulinemia. *J. Clin. Microbiol.* 2011;49, 3432 – 3435.
- [10]Sharp S E. Chronic Skin Lesions From a Patient With Bruton’s X-Linked Agammaglobulinemia. *J. Clin. Microbiol.* 2011;49, 483 – 48+.
- [11]Dua J, Elliot E, Bright P, et al. Pyoderma Gangrenosum-Like Ulcer Caused by *Helicobacter Cinaedi* in a Patient With X-Linked Agammaglobulinaemia. *Clin. Exp. Dermatol.* 2012;37, 642 – 645
- [12]Turvey SE, Leo SH, Boos A, et al. Successful Approach to Treatment of *Helicobacter Bilis* Infection in X Linked Agammaglobulinemia. *J. Clin. Immunol.* 2012;32, 1404 – 1408.

- [13]Toyofuku M, Tomida J., Kawamura Y,et al. Helicobacter Cinaedi Bacteremia Resulting From Antimicrobial Resistance Acquired During Treatment for X-Linked Agammaglobulinemia.*Infect. Chemother.* 2016;22, 704 – 706.
- [14]Degand N, Dautremer J, Pilmis B, et al. Helicobacter Bilis-Associated Suppurative Cholangitis in a Patient With X-Linked Agammaglobulinemia. *J. Clin. Immunol.* 2017;37, 727 – 731.
- [15]Hill A, Byrne A, Bouffard D,et al. Helicobacter Cinaedi Bacteremia Mimicking Eosinophilic Fasciitis in a Patient With X-Linked Agammaglobulinemia. *JAAD Case Rep.* 2018;4, 327 – 329.
- [16]Matsumoto A, Yeh I, Schwartz B,et al. Chronic Helicobacter Cinaedi Cellulitis Diagnosed by Microbial PolymeraseChain Reaction,2017; *JAAD Case Rep.* 3, 398 – 400.
- [17]Sugimoto M, Takeichi T, Muramatsu H, et al. Recurrent Cellulitis Caused by Helicobacter Cinaedi in a Patient With X-Linked Agammaglobulinaemia. *Acta Derm. Venereol.* 2017;97, 277 – 278.
- [18]Inoue K, Sasaki S, Yasumi T, et al. Helicobacter Cinaedi-Associated Refractory Cellulitis in Patients With XLinked Agammaglobulinemia. *J Clin Immunol* 2020;40, 1132 – 1137.
- [19]Fettinger NB, Bradley, JS, Ramchandar N. An 11-Year-old Male With X-linked Agammaglobulinemia and Persistent Abdominal Pain.*Pediatr Infect Dis J.*2021;1;40(4):356 -358
- [20]Romo-Gonzalez C , Bustamante-Ogando JC, Yamazaki-Nakashimada MA,et al.Infections With Enterohepatic Non- H. pylori Helicobacter Species in X-Linked Agammaglobulinemia: Clinical Cases and Review of the Literature.*Front Cell Infect Microbiol.* 2022;4:11:807136.
- [21]Lenskaya V , O'Connor M,de Moll EH,et al.Relapsing Helicobacter cinaedi Cellulitis without Bacteremia in a Patient with X-linked Agammaglobulinemia after 1 Year of Antibiotic Prophylaxis.*Skinme.* 2022 ;20;20(6):457-459.
- [22]Fjordside L, Herløv C ,Drabe CH,et al.Helicobacter trogontum Bacteremia and Lower Limb SkinLesion in a Patient with X-Linked Agammaglobulinemia-ACase Report and Review of the Literature.*Pathogens.*2022 ;27;11(11):1247.
- [23]Zhang L, Zhou ML, Lv W,et al.Metagenomics assists in the diagnosis of a refractory, culture-negative pyoderma gangrenosum-like ulcer caused by Helicobacter cinaedi in a patient with primary agammaglobulinemia.*J Microbiol Immunol Infect.*2023;56(6):1284-1287.