Article title: Patient and physician perceptions of painful diabetic peripheral neuropathy in **South-East Asia** 

Journal name: Advances in Therapy

Author names: Rayaz A. Malik, Emre Aldinc, Siew-Pheng Chan, Chaicharn

Deerochanawong, Chii-Min Hwu, Raymond L. Rosales, Chun-Yip Yeung, Koichi Fujii, and

**Bruce Parsons** 

Email of corresponding author: <a href="mailto:emre.aldinc@pfizer.com">emre.aldinc@pfizer.com</a>

Affiliation of corresponding author: Pfizer Inc, New York, NY, USA

1

### PHYSICIAN QUESTIONNAIRE

#### INTRODUCTION ONLINE

Thank you for agreeing to participate in this study. This interview is being conducted by Kantar Health, an independent market research agency to investigate doctors' attitudes towards treatment of diabetes in Asia.

The interview will last approximately 20 minutes. Your responses will be kept strictly confidential and your contact information will not be included in any market research report. Your opinions are important to us and would be very much appreciated. Please note that this interview is being carried out strictly for market research purposes and is in no way intended as a promotional exercise for any third party.

Kantar Health is abided by the Market Research Society guidelines and relevant data protection regulations. In line with these, you should be aware that:

- This Market Research is sponsored by a pharmaceutical company, this research study
  is not designed to be promotional in anyway, you have the right to withdraw from the
  interview at any time and withhold any information.
- All information shared with you in this interview should be kept confidential and should be assumed to represent hypotheses about what can or cannot be said about a product. It should not be used to influence any thinking or decisions taken outside this research setting.
- Please note that should you raise during this survey an adverse event in a specific
  patient of a specific drug, you will be obligated to report the adverse event directly to
  the Company's pharmaco-vigilance team or regulatory authorities.

This survey is being conducted on behalf of a pharmaceutical company.

# Please click on the button below to continue

This survey may contain confidential information. Any information obtained through completing this market research is confidential and must not be disclosed to third parties. Do you agree to keep this information confidential?

| Yes | 1 | Continue  |  |
|-----|---|-----------|--|
| No  | 2 | TERMINATE |  |

## **SECTION 1: PATIENT LOAD AND DIABETES MANAGEMENT PRIORITY**

| Q1.   | In a typical month, how many patients, of any condition, do you personally treat? |   |                   |   | ?   |
|---|---|---|-------------------|---|-----|
|   |   | Patients of any condition                               | on in a typical r | month   |     |
| Q2. Of the [pipe in answer from Q1] patients you personally treat in a typical remany do you treat with the following conditions? |   |   |                   |   | how |
|   |   | Conditions  |                   | Number of patients treated in a typical month |     |
|   |   | Diabetes (Type 1 and / or 2)                            | 1                 |   |     |
|   |   | Any other condition (all other non-diabetes conditions) | 2                 |   |     |
|   |   |   | TOTAL             | =Q1   |     |

**Q3.** Of those [pipe in response from  $Q2 = code\ 1$ ] diabetes patients that you treat in a typical month, what proportion suffers from the following conditions?

|   |       | Percentage (%) of patients |
|---|-------|----------------------------|
| Chronic heart disease                         | 1     | %                          |
| Obesity                                       | 2     | %                          |
| Hypertension                                  | 3     | %                          |
| Arthritis                                     | 4     | %                          |
| Painful diabetic peripheral neuropathy (pDPN) | 5     | %                          |
| Kidney disease                                | 6     | %                          |
|   | TOTAL | ≥ 100%                     |

Q4. What are your priorities when managing your diabetes patients? On a scale of 1–7, what priority would you put on the following when managing your diabetes patients (1 = lowest priority, 7= highest priority)?

|                          |   | Rating 1-7 |
|--------------------------|---|------------|
| Control of HbA1c         | 1 |            |
| Relief of pain from DPN  | 2 |            |
| Manage lipid disorder    | 3 |            |
| Maintain kidney function | 4 |            |

| Q5. | Based on your perception of the wider [pipe in country] market, approximately what |
|-----|--|
|     | proportion of diabetes patients suffer from painful DPN? [WHOLE NUMBER ONLY        |
|     | (allow from 1 to 100)]   |
|     | %  |

# **SECTION 2: IMPACT OF pDPN**

Q6. What is the impact of Painful DPN on patient's quality of life?

| ←Negligible Very Serious→ |   |   |   |   | ry Serious→ |   |
|---------------------------|---|---|---|---|-------------|---|
| 1                         | 2 | 3 | 4 | 5 | 6           | 7 |
|                           |   |   |   |   |             |   |

Q7. Among your patients with painful DPN, approximately what proportion are impacted by the following as a result of their painful DPN? [WHOLE NUMBER ONLY (allow from 0 to 100)]

| Sleep problems                                     | 1  | %         |
|--|----|-----------|
| Reduced ability to carry out normal duties at work | 2  | %         |
| Reduced ability to perform exercise                | 3  | %         |
| Depression   | 4  | %         |
| Anxiety  | 5  | %         |
| Poor appetite                                      | 6  | %         |
| Reduced libido                                     | 7  | %         |
| Increased cases of anger (i.e. due to frustration) | 8  | %         |
| Decreased desire to interact with others           | 9  | %         |
| Increased difficulty in holding concentration      | 10 | <b></b> % |

## **SECTION 3: DIABETIC PATIENT ENGAGEMENT AND DISCUSSION ON PAIN**

Q8. When discussing pain with your diabetes patients what proportion of the time do you initiate the conversation? How about the patient? [WHOLE NUMBER ONLY (allow from 0 to 100)]

| You, the doctor | 1     | %      |  |
|-----------------|-------|--------|--|
| The patient     | 2     | %      |  |
|                 | TOTAL | 100% % |  |

| Q9. | Considering all the diabetes patients you see in a typical month, what proportion do you |
|-----|--|
|     | screen for painful DPN? [WHOLE NUMBER ONLY (allow from 0 to 100)]                        |

|  | % |
|--|---|
|--|---|

Q10. Do you discuss pain with your painful DPN patients every time they visit you? [SINGLE RESPONSE ONLY]

| Yes | 1 |
|-----|---|
| No  | 2 |

Q11a. When discussing painful DPN with your diabetic patients, how do these patients typically describe the painful DPN they experience? Please review the following list. Please indicate which of the following words or terms patients use to describe their pain.

[MULTIPLE RESPONSES POSSIBLE]

Q11b. [Ask only if code 99 is not selected for Q11a] Please review the following list and indicate the order in terms of most commonly used words or terms patients have typically used to describe their painful DPN. Rank the order (max 3) of most commonly described pain symptoms: rank 1 for most common, rank 2 for second most common, and rank 3 for third most common.

|                                    |    | Q11a Please indicate which of the following words or terms patients use to describe their pain | Q11b<br>Rank<br>(Top 3 only) |
|------------------------------------|----|--|------------------------------|
| Burning sensation                  | 1  |  |                              |
| Numbing sensation                  | 2  |  |                              |
| Electric shocks                    | 3  |  |                              |
| Tingling pain                      | 4  |  |                              |
| Sharp or stabbing                  | 5  |  |                              |
| Pins & needles                     | 6  |  |                              |
| Dull ache                          | 7  |  |                              |
| Throbbing pain                     | 8  |  |                              |
| Other, please specify [open ended] |    |  |                              |
| None of these [single response]    | 99 |  |                              |

Q12a. How easy do you feel your patients find it to clearly describe their pain, without you prompting with terms or words?

| ←Very hard to describe Very easy to describe→ |   |   |   |   |   |   |
|---|---|---|---|---|---|---|
| 1   | 2 | 3 | 4 | 5 | 6 | 7 |
|   |   |   |   |   |   |   |

Q12b. How easy do you find it to interpret your patients' descriptions of their pain?

| ←Very hard to interpret Very easy to interpret |   |   |   |   | o interpret→ |   |
|--|---|---|---|---|--------------|---|
| 1  | 2 | 3 | 4 | 5 | 6            | 7 |
|  |   |   |   |   |              |   |

# SECTION 4: pDPN DIAGNOSIS, CLASSIFICATION, AND DISEASE MONITORING

Q13. Which of the following tools if provided would you find helpful to diagnose DPN?

[MULTIPLE RESPONSES ALLOWED]

| List of local language descriptions | 1 |
|-------------------------------------|---|
| Pain scale                          | 2 |
| Pain questionnaire                  | 3 |
| Others, please specify [open ended] |   |

Q14. For the diabetic patients that you see that do suffer from painful DPN, where do these patient mostly tell you their pain is coming from? Please indicate the site (or sites) of diabetes-related pain as voiced by your diabetic patients. [MULTIPLE RESPONSES POSSIBLE]

| Feet                      | 1 |
|---------------------------|---|
| Legs                      | 2 |
| Arm                       | 3 |
| Hands                     | 4 |
| Fingers and / or toes     | 5 |
| Others, please specify    |   |
| [open ended]              |   |
| Do not identify source of | 0 |
| pain [single response]    | 9 |

Q15. How long after the patient is diagnosed with diabetes do they develop painful DPN?

[SINGLE RESPONSE ONLY]

| Less than 1 year     | 1 |
|----------------------|---|
| Between 1 to 3 years | 2 |
| More than 3 years    | 3 |

**Q16a.** Of your patients that suffer from painful DPN that you see, what proportion fall into the following classification in terms of pain severity?

|   |     | Percentage of patients (%) |
|---|-----|----------------------------|
| 1 to 3 – mild pain (nagging, annoying, interfering with their daily activities)     | 1   | <b></b> %                  |
| 4 to 6 – moderate pain<br>(interferes significantly with<br>their daily activities) | 2   | <u></u> %                  |
| 7 to 10 – severe pain (disabling, unable to perform their daily activities)         | 3   | %                          |
|   | SUM | = 100%                     |

**Q16b.** In your opinion, how motivated are your diabetic patients in managing their pain condition compared to their existing concomitant conditions? Please rate how motivated you would consider the following patient groups (mild, moderate, severe) are to manage their diabetes related pain.

| ←Not motivated at all Extremely motivated→ |   |   |   |   |   |   |
|--|---|---|---|---|---|---|
| 1  | 2 | 3 | 4 | 5 | 6 | 7 |
|  |   |   |   |   |   |   |

**Q17.** Do you consider painful DPN a condition that requires a formal diagnosis? [SINGLE RESPONSE]

| Vac | 1             |  |  |
|-----|---------------|--|--|
| Yes | [Go to Q18]   |  |  |
| No  | 2             |  |  |
| No  | [Jump to Q21] |  |  |

Q18. Do you typically diagnose your diabetic patients with painful DPN yourself or is this done by someone else? Who typically diagnoses your diabetic patients for painful DPN?

[SINGLE RESPONSE]

| I typically perform the diagnosis myself         | 1 |
|--|---|
| Cardiologists                                    | 2 |
| Endocrinologists / diabetologists                | 3 |
| Pain specialists                                 | 4 |
| Nephrologist                                     | 5 |
| Orthopedic surgeons                              | 6 |
| General practitioner / primary care practitioner | 7 |
| Others, please specify [open ended]              | 8 |

Q19. [Ask if Q18 = code 1] How do you personally diagnose painful DPN? [MULTIPLE RESPONSES POSSIBLE]

| Clinical examination   | 1 |
|------------------------|---|
| Symptom observation    | 2 |
| Nerve conduction test  | 3 |
| Foot examination       | 4 |
| Monofilament test      | 5 |
| Medical history        | 6 |
| Risk factors, please   | 7 |
| specify [open ended]   | 7 |
| Others, please specify |   |
| [open ended]           |   |

**Q20.** [Ask if Q18 = code 1] For how long do you monitor the condition of your patients that suffer from Painful DPN? [SINGLE RESPONSE]

| <1 month   | 1 |
|------------|---|
| 1–3 months | 2 |

| > 3 months up to 1 year | 3 |
|-------------------------|---|
| >1 year up to 2 years   | 4 |
| >2 years                | 5 |

## SECTION 5: TREATMENT, BARRIERS, AND THE USE OF PAIN GUIDELINES

Q21a. In your opinion, what are the key barriers to treatment when managing painful DPN patients? Please review the following list and indicate whether each are key barriers to treatment when managing diabetes-related pain. Please answer only for your total diabetes-related pain patients. [MULTIPLE RESPONSES POSSIBLE]

**Q21b.** Please review the same list and indicate the order of most common barriers. Rank the order (max 3) of most common barrier. Rank 1 for most common, rank 2 for second most common, and rank 3 for third most common.

|  | Q21a Please indicate which of the following are key barriers to treatment | Q21b<br>Rank<br>(Top 3 only) |
|--|---|------------------------------|
| Patients do not initiate a discussion on their painful DPN | 1   |                              |
| Patients do not like to discuss their painful DPN          | 2   |                              |
| Patients are fearful of additional costs of treatment      | 3   |                              |
| Patients lack awareness of diabetes specific related pain  | 4   |                              |
| Patients find it difficult to describe their pain          | 5   |                              |
| Patients refuse to accept the seriousness of painful DPN   | 6   |                              |
| Painful DPN is viewed to be unimportant to manage          | 7   |                              |
| Limited consultation time                                  | 8   |                              |
| Others, please specify [open ended]                        |   |                              |

**Q22.** For those diabetic patients that are diagnosed with painful DPN, who typically manages it? [SINGLE RESPONSE]

| Myself   | 1 |
|--|---|
| Cardiologists                                    | 2 |
| Endocrinologists / diabetologists                | 3 |
| Pain specialists                                 | 4 |
| Nephrologist                                     | 5 |
| Orthopedic surgeons                              | 6 |
| General practitioner / primary care practitioner | 7 |
| Others, please specify [open ended]              | 8 |

**Q23.** Which of the following (if any) do you typically recommend to patients suffering from mild, moderate and severe DPN? [MULTIPLE RESPONSES POSSIBLE FOR EACH TYPE OF PATIENTS]

|  |   | Patient types (by pain severity)                                       |  |  |
|--|---|--|--|--|
|  |   | Mild pain (nagging, annoying, interfering with their daily activities) | Moderate pain (interferes significantly with their daily activities) | Severe pain<br>(disabling,<br>unable to<br>perform their<br>daily<br>activities) |
| Change of lifestyle (diet and exercise change) to ease symptoms of pain                                    | 1 |  |  |  |
| Prescription medication to ease symptoms of pain   | 2 |  |  |  |
| Traditional medication (e.g. herbs, Chinese medication) to ease symptoms of pain                           | 3 |  |  |  |
| Vitamin treatment and health supplements to ease symptoms of pain  | 4 |  |  |  |
| Better regulate existing diabetes treatment (i.e. tightening blood sugar control) to ease symptoms of pain | 5 |  |  |  |
| Other, please specify [open ended]   | 6 |  |  |  |

**Q24a.** What diabetes guidelines, if any, do you adhere to in the management of your diabetes patients? [MULTIPLE RESPONSES POSSIBLE]

**Q24b.** Are you aware if these guidelines have a section on the management of painful DPN? [MULTIPLE RESPONSES POSSIBLE]

|  |   | Q24a | Q24b |
|--|---|------|------|
| American Diabetes Association (ADA)  | 1 |      |      |
| International Diabetes Federation (IDF)  | 2 |      |      |
| Archives of Physical Medicine and Rehabilitation (APMR)  | 3 |      |      |
| American Academy of Neurology (AAN)  | 4 |      |      |
| European Association for the Study of Diabetes (EASD)  | 5 |      |      |
| Local Guidelines   | 6 |      |      |
| I am not aware that these guidelines have a section on the management of diabetes related pain | 7 |      |      |
| Others, please specify [open ended]  |   |      |      |

THANK YOU AND CLOSE.