

Electronic supplementary material

Table 1 Baseline characteristics of FHBL patients and controls

Characteristic	FHBL (n=7)	Control (n=7)	p value
Age (years)	47 (22–61) 44±5.6	45 (22–60) 42±5	NS
BMI (kg/m ²)	27 (22–32) 27±1.2	26 (20–36) 27±1.8	NS
WHR (cm)	1.02 (0.97–1.05) 1.02±0.01	1.01 (0.94–1.12) 1.02±0.03	NS
Total cholesterol (mmol/l)	1.94 (1.39–2.81) 1.97±0.18	4.75 (3.47–5.23) 4.49±0.22	<0.01
Triacylglycerol (mmol/l)	0.14 (0.05–0.49) 0.20±0.06	0.76 (0.40–1.07) 0.73±0.08	<0.01
ApoB (g/l)	0.24 (0.12–0.4) 0.24±0.04	0.83 (0.54–0.90) 0.78±0.05	<0.01
HDL (mmol/l)	1.19 (0.7–1.53) 1.14±0.11	1.3 (0.79–1.83) 1.3±1.15	NS
HOMA index	1.58 (0.21–2.35) 1.50±0.25	0.76 (0.23–1.58) 0.84±0.20	NS
Aspartate aminotransferase (U/l)	39 (22–49) 37±4	28 (14–34) 25±3	<0.05
Alanine aminotransferase (U/l)	62 (22–106) 63±13	23 (10–32) 21±3	0.05
C-reactive protein (mg/l)	1.4 (0.5–3.1) 1.6±0.4	1.0 (0.5–6.2) 1.6±0.8	NS

Data are median (range) and mean±SEM