

## Electronic supplementary material

**Table 2** Body fat composition

Variable	FHBL ( <i>n</i> =7)	Controls ( <i>n</i> =7)	<i>p</i> value
Body weight (kg)	86 (70–110)	87 (71–114)	NS
	88±5	90±5	
Body fat (%)	26 (19–30)	23 (12–35)	NS
	25±2	24±3	
Lean body mass (kg)	61 (49–77)	64 (58–67)	NS
	61±39	63±12	
Peripheral fat mass (kg)	8.4 (5.7–14.1)	8.6 (3.9–16.6)	NS
	8.8±10.3	9.5±15.7	
Trunk fat mass (kg)	11.7 (6.1–16.7)	10.8 (3.6–19.9)	NS
	11.8±13.1	11.0±18.4	
Abdomen			
Subcutaneous adipose tissue (cm <sup>2</sup> )	189 (148–429)	203 (48–505)	NS
	233±404	236±531	
Visceral adipose tissue (cm <sup>2</sup> )	124 (39–181)	107 (27–219)	NS
	128±175	115±237	
Visceral:subcutaneous adipose tissue ratio	0.62 (0.26–1.19)	0.55 (0.22–1.08)	NS
	0.63±0.14	0.56±0.11	

Data are median (range) and mean±SEM