

Manuscript Title: Changes in inflammatory factors in the Brown Norway rat model of food allergy

Qingling Zhu, PhD^{a,b}, Junli Wang, PhD^a, Jingqiu Ma, PhD^a, Xiaoyang Sheng, PhD^{a*}, Feng Li, PhD^{a*}

^a Department of Child and Adolescent Healthcare, MOE-Shanghai Key Laboratory of Children's Environmental Health, Xinhua Hospital Affiliated to Shanghai Jiao Tong University School of Medicine, Shanghai, China 200092 ;

^b Department of Children Healthcare, Quanzhou Women's and Children's Hospital, Quanzhou, Fujian, China 362000

***Corresponding author:**


1. Feng Li, Department of Child and Adolescent Healthcare, Xinhua Hospital Affiliated to Shanghai Jiao Tong University School of Medicine, No. 1665 Kongjiang Road, Yangpu Shanghai, China, 200092
Tel: +86- 13512111965, E-mail: lifeng@xinhumed.com.cn

2. Xiaoyang Sheng, Department of Child and Adolescent Healthcare, Xinhua Hospital Affiliated to Shanghai Jiao Tong University School of Medicine, No. 1665 Kongjiang Road, Yangpu Shanghai, China, 200092


Tel: +86-13818072645, E-mail: shengxiaoyangcn@aliyun.com

Supplementary Info File:


Multiple exposures of Weston blot in Fig 3



Sup 1 The levels of S100A8/A9 in the experimental group




Sup 2 The levels of S100A8/A9 in the control group



Sup 3 The levels of TLR4 in the experimental group



Sup 4 The levels of TLR4 in the control group



Sup 5 The levels of TNF- α in the experimental group



Sup 6 The levels of TNF- α in the control group



Sup 7 The levels of GAPDH in the experimental group



Sup 8 The levels of GAPDH in the control group