

Additional file 3. Risk factor values for CKD patients who had above-target values at baseline

	Intervention	Control	p-value
LDL-C \geq2.5/1.8[†] baseline, n	54	52	
LDL-C baseline, median (25th–75th percentile)	3.0 (2.7-3.5)	3.1 (2.7-3.7)	0.33
LDL-C 12 months, median (25th–75th percentile)	2.2 (1.8-2.8) ^{***}	3.0 (2.6-3.7) ^{NS}	<0.001
SBP \geq140 baseline, n	47	54	
SBP baseline, median (25th–75th percentile)	150 (144-155)	150 (140-160)	0.96
SBP 12 months, median (25th–75th percentile)	132 (126-150) ^{***}	145 (130-160) ^{NS}	0.05
DBP \geq90 baseline, n	12	14	
DBP baseline, median (25th–75th percentile)	92 (90-99)	90 (90-94)	0.37
DBP 12 months, median (25th–75th percentile)	80 (71-84) ^{**}	84 (79-94) ^{**}	0.07

^{**}p \leq 0.01, ^{***}p \leq 0.001, indicating a significant change of median values within each group between baseline and 12 months; ; [†]The target LDL-C value was <2.5 mmol/L until 31 March, 2013, when local guidelines for diabetic patients changed this target to <1.8 mmol/L. LDL-C: low density lipoprotein cholesterol (mmol/L); n: number of valid cases; SBP: systolic blood pressure (mmHg); DBP: diastolic blood pressure (mmHg); NS: non-significant.