ADDITIONAL FILE 2

Analyses of exercise time (measured as mean difference, and subgroup analyses)


CCS class change and anginal episodes

The sham effect of invasive interventions in chronic coronary syndromes – a systematic review and meta-analysis

Authors: Catarina Palma, Cláudio David MD, Ricardo M Fernandes MD PhD, Fausto J Pinto MD PhD, João Costa MD PhD, Joaquim J Ferreira MD PhD, Daniel Caldeira MD PhD


Table of contents

Supplementary Figure: Mean difference of exercise time
Supplementary Figure: Analysis of relative change for exercise time according with the type of sham
Supplementary Figure: Analysis of relative change for exercise time according with the mean LVEF/proportion of HF patients. HF: Heart failure; NR: not reported
Supplementary Figure: Mean difference of Canadian Cardiovascular Society anging class.
Supplementary Figure: Mean difference of anginal episodes and nitroglycerine use per week.




Supplementary Figure: Mean difference of exercise time

Pooled and individual mean difference and 95% CI for the exercise time outcome with Bruce protocol, modified Bruce protocol and bicycle protocol.




Supplementary Figure: Analysis of relative change for exercise time according with the type of sham

Pooled and individual relative effect and 95% CI for exercise time according with the type of sham (pure sham intervention or sham intervention + placebo solution).




Supplementary Figure: Analysis of relative change for exercise time according with the mean LVEF/proportion of HF patients. HF: Heart failure; NR: not reported



Supplementary Figure: Mean difference of Canadian Cardiovascular Society angina class.

Pooled and individual mean difference and 95% CI for the CCS angina class outcome.



Supplementary Figure: Mean difference of anginal episodes and nitroglycerine use per week.

Pooled and individual mean difference and 95% CI for the anginal episodes and nitroglycerine use per week outcome.