

Table 2. Sensitivity of the bacteria isolated from jejunal aspirate to various antibiotics.

Bacteria	Sensitive to							
	Ampicillin	Ampicillin Sulbactam	Tetracyclin	Cotrimox	Erythro	Norflox	Ciproflo	Cefalexin
<i>E. coli</i> (n = 12)	3	5	4	5	NT	7	9	5
<i>Streptococcus sp.</i> (n = 12)	9	11	6	5	9	11	12	9
<i>E. fecalis</i> (n = 3)	3	3	1	0	1	1	1	NT
<i>E. fecium</i> (n = 1)	1	1	0	0	1	0	0	NT
<i>S. aureus</i> (n = 5)	3	4	2	5	3	4	5	3
<i>C. freundii</i> (n=1)	0	1	0	1	NT	1	1	1
<i>K. pneumoniae</i> (n = 5)	1	3	2	3	NT	3	4	3
<i>P. mirabilis</i> (n =1)	1	1	0	0	NT	1	1	1
<i>A. baumannii</i> (n = 3)	0	0	2	2	NT	3	3	3
<i>P. aeruginosa</i> (n = 3)	NT	NT	1	2	NT	3	3	NT
<i>B. melaninogenicus</i> (n = 1)	0	1	1	1	0	0	0	0
Total (n = 47)	21/44 (48)	30/44 [*] (68)	19/47 ^{*, \$} (40)	24/47 (51)	14/22 (64)	34/47 [*] (72)	39/47 ^{\$} (83)	25/40 (62.5)

Figures within parenthesis indicate percentages. NT: not tested.

*p < 0.01; \$p <0.0001 (Chi-squared test)