

Table 4 The adjusted odds ratios of metabolic risk factors for NAFLD

Variable(s)	Male				Female			
	β	S.E.	OR	<i>P</i> -value	β	S.E.	OR	<i>P</i> -value
BMI	1.195	.014	3.304	.000	1.336	.023	3.806	.000
WC	.673	.014	1.960	.000	1.046	.022	2.846	.000
BP	.667	.011	1.949	.000	0.719	.017	2.053	.000
FPG	.407	.015	1.502	.000	0.899	.024	2.458	.000
TG	1.337	.012	3.809	.000	1.218	.020	3.381	.000
HDL	.468	.021	1.596	.000	0.588	.024	1.801	.000
normal	-6.239	.036	0.002	.000	-8.096	.048	0.000	.000

β :Beta coefficient; S.E: Standard error; OR: odds ratio; BMI: Body mass index, $\geq 25\text{kg/m}^2$; WC: waist circumference, $\geq 90\text{cm}$ (male), $\geq 80\text{cm}$ (female);

BP: blood pressure, $\geq 130/85\text{mmHg}$ Or hypertension (HBP); FPG: Fasting blood glucose, $\geq 5.6\text{ mmol/L}$ Or type 2 diabetes (2-DM);

TG: Triglycerides, $\geq 1.7\text{mmol/L}$; HDL: High Density Lipoprotein, $\leq 1.03\text{mmol/L}$, (male) $< 1.29\text{mmol/L}$ (female).