

	Instruments	When & who?	Baseline	3 months (= ca. 12-13 weeks)	6 months (= ca. 26 weeks)
<i>Selection criteria</i>	- GDS (Reisberg) - MDS-CPS	Pre selection Pre selection	NHP NHP		
<i>Primary Outcome</i>	- CMAI - MDS-RAI pain scale - NPI-NH - PACSLAC-D	Incl. Incl. Incl. Incl.	Nurse Nurse Nurse Nurse & VIDEO	Nurse Nurse Nurse Nurse & VIDEO	Nurse Nurse Nurse Nurse
<i>Secondary Outcome</i>	- Qualidem - Cornell & MDS-DRS - Medication - DS-DAT - FACS	Incl. Incl. Incl. Incl. Incl.	Nurse Nurse RA VIDEO VIDEO	Nurse Nurse RA VIDEO VIDEO	Nurse Nurse RA
<i>Control variables / Demographics</i>	- (co-) Morbidity - Hospitalisation, fixation etc. - Demographics; sex, age, background - ADL (Katz) - Apolipoprotein E4	Incl. Incl. Incl. Incl. Incl.	NHP RA RA Nurse RA	RA	RA

NHP = Nursing Home Physician, Nurse = caregiver, RA = research assistant, Incl. = included residents