
Additional file 1. Predictors of regular dizziness in an older community population at 7- and 10-year follow-up – Additional analysis: *without* 'history of dizziness' as potential predictor

	B	SE	Wald	OR	95% CI	P value
Characteristic						
<i>7-year follow-up</i> ^a						
Years of education	-0.066	0.036	3.3	0.94	0.9-1.0	0.069
Living alone	0.422	0.223	3.6	1.5	0.9-2.4	0.058
History of osteo/rheumatoid arthritis	0.762	0.233	10.7	2.1	1.4-3.4	0.001
Use of nitrates	0.879	0.342	6.6	2.4	1.2-4.7	0.010
Presence of anxiety or depression	0.735	0.247	8.9	2.1	1.3-3.4	0.003
Impaired vision	0.698	0.245	8.1	2.0	1.2-3.2	0.004
Impaired function of lower extremities (chair stands)	0.704	0.222	10.0	2.0	1.3-3.1	0.002
<i>10-year follow-up</i> ^b						
Living alone	0.466	0.269	3.0	1.6	0.9-2.7	0.083
History of cancer	0.568	0.348	2.7	1.8	0.9-3.5	0.103
Use of anxiolytics	0.902	0.453	4.0	2.5	1.0-6.0	0.046
Impaired vision	0.497	0.314	2.5	1.6	0.9-3.0	0.114
Impaired function of lower extremities (chair stands)	0.894	0.271	10.9	2.4	1.5-4.2	0.001

CI: Confidence Interval; OR: Odds Ratio; SE: Standard Error.

*: Multivariable logistic regression analysis, using backward elimination (Wald test) with a *P* value of 0.157 used for removal.

^a: AUC 0.75 [0.70-0.80].

^b: AUC 0.68 [0.61-0.75].