Additional file 1. Frailty Index Post-Acute Care

	icode	Question	Value Labels	InterRAI Cut Offs	FI score	Running Denominator
	iC1	Cognitive skills for daily decision making	0 Independent	0	0	1
			1 Modified independence			•
			2 Minimally impaired	1 – 2	0.5	
			3 Moderately impaired			
			4 Severely impaired	3 - 5	1	
			5 No discernible consciousness, coma			
	iC2a	Short term memory	0 Memory OK	0	0	2
	1000	,	1 Memory problem	1	1	_
	iC2b	Long term memory	0 Memory OK	0	0	3
	1025	Long tom momory	1 Memory problem	1	1	
	iC2c	Procedural memory	0 Memory OK	0	0	4
_	1020	1 recodular memory	1 Memory problem	1	1	7
fjo	iC2d	Situational memory	0 Memory OK	0	0	5
gni	1024	Oldational memory	1 Memory problem	1	1	
C. Cognition	iC3a	Easily distracted	Behaviour not present	0	0	6
ပ	103a	Lasily distracted	Behaviour present, consistent with usual functioning			
			2 Behaviour present, different from usual functioning	1 - 2	1	
	iC3b	Disorganised speech	Dehaviour not present	0	0	7
	1036	Disorganised speech	Behaviour present, consistent with usual functioning			,
			2 Behaviour present, different from usual functioning	1 - 2	1	
	iC3c	Mental function varies over course of day	Dehaviour not present	0	0	8
	1030	Werital function values over course of day	Behaviour present, consistent with usual functioning			
			2 Behaviour present, different from usual functioning	1 - 2	1	
			2 Benaviour present, different from usual functioning	1-2	'	
	iC4	Acute change in mental status	0 No	0	0	9
	104	Acute change in mental status	1 Yes	1 1	1	9
			1 165	'	'	
	iD3	Hearing	0 Adequate	0	0	10
ion			1 Minimal difficulty			
\ \ Si			2 Moderate difficulty	1	0.5	
p			3 Severe difficulty			
la (4 No hearing	2 - 4	1	
Communication and Vision	iD5	Vision	0 Adequate	0	0	11
ica			1 Minimal difficulty			
חשר			2 Moderate difficulty	1	0.5	
ШŢ			3 Severe difficulty			
ပိ			4 No vision	2 - 4	1	
<u> </u>						
0.0						

	icode	Question	Value Labels	InterRAI Cut Offs	FI score	Running Denominator
d and iour	iE1a-g	Indicators of depression, anxiety or sad mood Depression Rating Scale (0-14)	Calculate the Depression Rating Scale	DRS 0 DRS 1-2 DRS ≥3	0 0.5 1	12
E. Mood and Behaviour	iE3b- d,f	Indicators of behaviour symptoms Aggressive Behaviour Scale (0-12)	Calculate the Aggressive Behaviour Scale	ABS 0 ABS 1-3 ABS ≥4	0 0.5 1	13
	iF1a	Bathing	0 Independent 1 Independent, setup help only 2 Supervision- oversight, cueing	0 - 1 2	0 0.5	14
			3 Limited assistance- guided manoeuvring of limbs 4 Extensive assistance- 1 person weight bearing support 5 Maximal assistance- 2 person weight bearing support 6 Total dependence 8 Activity did not occur	3 – 6, 8	1	
	iF1b	Personal hygiene	0 Independent 1 Independent, setup help only 2 Supervision- oversight, cueing 3 Limited assistance- guided manoeuvring of limbs	0–1 2	0 0.5	15
atus			Extensive assistance- 1 person weight bearing support Maximal assistance- 2 person weight bearing support Total dependence Activity did not occur	3 – 6, 8	1	
F. Functional status	iF1c	Dressing upper body	1 Independent 1 Independent, setup help only 2 Supervision- oversight, cueing 3 Limited assistance- guided manoeuvring of limbs	0 - 1 2	0 0.5	16
<u>г.</u>			4 Extensive assistance- 1 person weight bearing support 5 Maximal assistance- 2 person weight bearing support 6 Total dependence 8 Activity did not occur	3 – 6, 8	1	
	iF1d	Dressing lower body	Independent Independent, setup help only Supervision- oversight, cueing Limited assistance- guided manoeuvring of limbs	0 - 1 2	0.5	17
			4 Extensive assistance- 1 person weight bearing support 5 Maximal assistance- 2 person weight bearing support 6 Total dependence 8 Activity did not occur	3 – 6, 8	1	

	icode	Question	Value Labels	InterRAI Cut Offs	FI score	Running Denominator
	iG1	Bladder continence	Continent Control with catheter or ostomy	0 – 1	0	33
			2 Infrequently incontinent 3 Occasionally incontinent	2 – 3	0.5	
			4 Frequently incontinent 5 Incontinent 8 Did not occur – no urine output in last 3 days	4,5,8	1	
	iF1g	Toilet Transfers	0 Independent	0 – 1	0	19
			1 Independent, setup help only2 Supervision- oversight, cueing3 Limited assistance- guided manoeuvring of limbs	2	0.5	
			4 Extensive assistance- 1 person weight bearing support 5 Maximal assistance- 2 person weight bearing support 6 Total dependence 8 Activity did not occur	3 – 6, 8	1	
	iF1h	Toilet use	0 Independent 1 Independent, setup help only	0 – 1	0	20
nence			2 Supervision- oversight, cueing 3 Limited assistance- guided manoeuvring of limbs	2	0.5	
G. Continence			4 Extensive assistance- 1 person weight bearing support 5 Maximal assistance- 2 person weight bearing support 6 Total dependence 8 Activity did not occur	3 – 6, 8	1	
	iF1i	Bed mobility	0 Independent 1 Independent, setup help only	0 – 1	0	21
			2 Supervision- oversight, cueing 3 Limited assistance- guided manoeuvring of limbs	2	0.5	
			4 Extensive assistance- 1 person weight bearing support 5 Maximal assistance- 2 person weight bearing support 6 Total dependence 8 Activity did not occur	3 – 6, 8	1	
	iF1j	Eating	0 Independent 1 Independent, setup help only	0 – 1	0	22
			2 Supervision- oversight, cueing 3 Limited assistance- guided manoeuvring of limbs	2	0.5	
			4 Extensive assistance- 1 person weight bearing support 5 Maximal assistance- 2 person weight bearing support 6 Total dependence 8 Activity did not occur	3 – 6, 8	1	
			o Activity did Not occur			

	icode	Question	Value Labels	InterRAI Cut Offs	FI score	Running Denominator
	iF2a	Mode of locomotion	Walking, no assistive device Walking, uses assistive device Wheelchair, scooter Bedbound	0 1 2-3	0 0.5 1	23
	iF2b	Walking speed	Count Timed 4-meter walk distance (m)/ walking time (s)	>= 0,8 m/s < 0,8 m/s Not performed	0 1 1	24
	iG1	Bladder continence	0 Continent 1 Control with catheter or ostomy 2 Infrequently incontinent 3 Occasionally incontinent 4 Frequently incontinent 5 Incontinent 8 Did not occur – no urine output in last 3 days	0 – 1 2 – 3 4,5,8	0 0.5 1	25
	iG4	Bowel continence	0 Continent 1 Control with ostomy 2 Infrequently incontinent 3 Occasionally incontinent 4 Frequently incontinent 5 Incontinent 8 Did not occur – no bowel movement in last 3 days	0 – 1 2 – 3 4,5,8	0 0.5 1	26
	iH1a	Hip fracture during last 30 days	0 Not present 1 Primary diagnosis of current stay 2 Diagnosis present, receiving active treatment 3 Diagnosis present, monitored but no active treatment	0 1-3	0 1	27
Diagnoses	iH1b	Other fracture during last 30 days	Not present Primary diagnosis of current stay Diagnosis present, receiving active treatment Diagnosis present, monitored but no active treatment	0 1-3	0 1	28
H. Diag	iH1c	Alzheimers disease	Not present Primary diagnosis of current stay Diagnosis present, receiving active treatment Diagnosis present, monitored but no active treatment	0 1-3	0	29
	iH1d	Dementia other than ALZ	Not present Primary diagnosis of current stay Diagnosis present, receiving active treatment Diagnosis present, monitored but no active treatment	0 1-3	0	30

	icode	Question	Value Labels	InterRAI Cut Offs	FI score	
	iH1h	Parkinson's disease	Not present Primary diagnosis of current stay Diagnosis present, receiving active treatment Diagnosis present, monitored but no active treatment	0 1-3	0	31
	iH1j	Stroke/ SVA	Not present Primary diagnosis of current stay Diagnosis present, receiving active treatment Diagnosis present, monitored but no active treatment	0 1-3	0	32
	iH1k	Coronary heart disease	Not present Primary diagnosis of current stay Diagnosis present, receiving active treatment Diagnosis present, monitored but no active treatment	0 1-3	0 1	33
	iH1I	Congestive heart failure	Not present Primary diagnosis of current stay Diagnosis present, receiving active treatment Diagnosis present, monitored but no active treatment	0 1-3	0 1	34
	iH1m	COPD	Not present Primary diagnosis of current stay Diagnosis present, receiving active treatment Diagnosis present, monitored but no active treatment	0 1-3	0 1	35
	iH1n	Anxiety	Not present Primary diagnosis of current stay Diagnosis present, receiving active treatment Diagnosis present, monitored but no active treatment	0 1-3	0 1	36
	iH1p	Depression	Not present Primary diagnosis of current stay Diagnosis present, receiving active treatment Diagnosis present, monitored but no active treatment	0 1-3	0 1	37
	iH1t	Cancer	Not present Primary diagnosis of current stay Diagnosis present, receiving active treatment Diagnosis present, monitored but no active treatment	0 1-3	0 1	38
	iH1u	Diabetes	Not present Primary diagnosis of current stay Diagnosis present, receiving active treatment Diagnosis present, monitored but no active treatment	0 1-3	0	39
I. Health Conditions	il1	Falls	0 No fall in last 90 days 1 No fall in last 30 days but fell 31-90 days ago 2 One fall in last 30 days 3 Two or more falls in last 30 days	0 1 2, 3	0 0.5 1	40

icode	Question	Value Labels	InterRAI Cut Offs	FI score	Running Denominator
il3a	Balance: Difficult to move self to standing	Not present Present but not exhibited last 3 days	0	0	41
		2 Exhibited on 1 of last 3 days 3 Exhibited on 2 of last 3 days 4 Exhibited daily in last 3 days	1-4	1	
il3b	Balance: Difficult to turn self around	Not present Present but not exhibited last 3 days Exhibited on 1 of last 3 days	0	0	42
		3 Exhibited on 2 of last 3 days 4 Exhibited daily in last 3 days	1-4	'	
il3c	Dizziness	Not present Present but not exhibited last 3 days	0	0	43
		2 Exhibited on 1 of last 3 days 3 Exhibited on 2 of last 3 days 4 Exhibited daily in last 3 days	1-4	1	
il3d	Unsteady gait	Not present Present but not exhibited last 3 days	0	0	44
		2 Exhibited on 1 of last 3 days 3 Exhibited on 2 of last 3 days 4 Exhibited daily in last 3 days	1-4	1	
i3l	Constipation	Not present Present but not exhibited last 3 days	0	0	45
		2 Exhibited on 1 of last 3 days 3 Exhibited on 2 of last 3 days 4 Exhibited daily in last 3 days	1-4	1	
il3o	Difficulty falling asleep	Not present Present but not exhibited last 3 days	0	0	46
		2 Exhibited on 1 of last 3 days 3 Exhibited on 2 of last 3 days 4 Exhibited daily in last 3 days	1-4	1	
il4	Dyspnoea	O Absence of symptom Absent at rest, but present with moderate activity	0	0	47
		Absent at rest, but present with normal activity Present at rest	1-3	1	
il5	Fatigue- inability to complete normal activity	0 None 1 Minimal	0	0	48
		2 Moderate 3 Severe	1	0.5	
		4 Unable to commence any day-to-day activity	2 - 4	1	

	icode	Question	Value Labels	InterRAI Cut Offs	FI score	Running Denominator
	i6a-b	Indicators of pain PAIN-scale (0-4)	Calculate the PAIN-scale	PAIN 0 PAIN 1 PAIN 2-4	0 0.5 1	49
	il7c	End-stage disease	0 No 1 Yes	0 1	0 1	50
	iJ1ab, iJ1bb	Calculate BMI	BMI	20 - 30 < 20 or >30	0	51
sr	iJ2a	Weight loss of 5% or more in last 30 days, 10% or more in last 180 days	0 No 1 Yes	0 1	0	52
onal Statı	iJ2b	Dehydrated	0 No 1 Yes	0 1	0 1	53
J. Oral and Nutritional Status	iJ3	Mode of intake	O Normal Modified independent –limited solid food Requires diet modification to swallow solid food Requires diet modification to swallow liquids Can swallow only pureed solids and thickened fluids Combined oral and parenteral or tube feeding Nasogastric tube feeding only	0 1-2 3 - 9	0 0.5	54
			7 Abdominal feeding tube e.g. PEG 8. Parenteral feeding only 9 Activity did not occur	3-3	'	
	iK1- iK2	Pressure ulcer-current or prior	0 No 1 Yes	0 1	0 1	55
Skin	iK3- iK6	Skin problems (any of presence of skin ulcer, major skin problem, skin tears, other skin conditions)	0 ≥1	0 1	0	56
Ϋ́	iK7	Foot problems	O No foot problems Foot problems, no limitation in walking Foot problems limit walking Foot problems prevent walking Foot problems, does not walk for other reasons	0 1-4	0 1	57

The denominator from PAC ITEMS is 57. For complete interRAI PAC forms, the individual's FI is calculated by dividing their number of deficits by 57. The denominator can be adjusted for missing items.