



Days until antiretroviral therapy was initiated

▲ bacteriologic-confirmed TB

● all patients

- - - fitted line (bacteriologic-confirmed TB)

— fitted line (all patients)

Days until ART was initiated	0-30 vs. >30	0-40 vs. >40	0-50 vs. >50	0-60 vs. >60	0-70 vs. >70	0-80 vs. >80	0-90 vs. >90	0-100 vs. >100	0-110 vs. >110	0-120 vs. >120
Log ₁₀ hazard ratio*	-0.15	-0.30	-0.15	0.08	0.04	0.08	0.08	0.20	0.30	0.36
95% Lower CI	-0.70	-1.00	-0.70	-0.52	-0.52	-0.52	-0.52	-0.40	-0.30	-0.22
95% Upper CI	0.46	0.23	0.40	0.65	0.61	0.66	0.67	0.82	0.91	0.97
Log ₁₀ hazard ratio†	-0.10	-0.40	-0.15	0.23	0.53	0.59	0.70	0.84	0.90	0.95
95% Lower CI	-1.00	-1.00	-1.00	-0.52	-0.40	-0.30	-0.22	-0.10	0.00	0.04
95% Upper CI	0.93	0.49	0.70	1.04	1.43	1.48	1.58	1.76	1.80	1.86