

Table 4: Distribution of PfCRT polymorphism according to the drug sensitivities

Drug	Profils	n	N° (%), of strains with polymorphism : (<i>PfCRT</i>)			
			HAPLOTYPE 72-76			p
			CVIEK	CVIET	MIX	
CQ	Resistant (IC50 ≥ 100 nM)	20	0 (0.0%)	20 (100.0%)	0 (0.0%)	0.56
	Sensitive (IC50 < 100 nM)	26	1 (3.8%)	25 (96.2%)	0 (0.0%)	
MDAQ	Resistant (IC50 ≥ 60 nM)	24	1 (4.2%)	23 (95.8%)	0 (0.0%)	0.29
	Sensitive (IC50 < 60 nM)	22	0 (0.0%)	22 (100%)	0 (0.0%)	
MF	Resistant (IC50 ≥ 20 nM)	11	0 (0.0%)	11 (100.0%)	0 (0.0%)	0.58
	Sensitive (IC50 < 20 nM)	35	1 (2.9%)	34 (97.1%)	0 (0.0%)	
DHA	Reduced susceptibility (IC50 ≥ 10 nM)	11	1 (9.1%)	10 (90.9%)	0 (0.0%)	0.23
	Sensitive (IC50 < 10 nM)	35	0 (0.0%)	35 (100.0%)	0 (0.0%)	