Appendix Table A4. Overview of studies included in the meta-analysis of spatial risk factors for tick-borne diseases and tick bites.

Citation	Cases	Controls	Age	Study dates	Outcome measure	Study type	Location
Aenishaenslin et al. 2017 [1]	42	2834	All	2014	disease	survey	Canada
Connally et al. 2009 [2]	364	349	All	2005- 2007	disease	case control	СТ
Mead et al. 2018[3]	91	840	All	2013	bite	case control	СТ
Vazquez et al. 2008 [4]	285	557	All	2003	disease	case control	СТ
Hinckley et al. 2015 bite [5]	1303	2502	All	2011- 2012	bite	controlled trial	CT, MD, NY
Hinckley et al. 2015 disease [5]	1273	2461	All	2011- 2012	disease	controlled trial	CT, MD, NY
Klein et al. 1996 [6]	42	42	Children	not specified	disease	case control	DE, MD, PA, NJ
Richards et al. 2015 [7]	22	13	Adults	2013	bite	controlled trial	KY, NC, OH, TN, VA, WV
Glass et al. 1995 [8]	47	492	not specified	1989- 1990	disease	case control	MD
Bowen et al. 1984 [9]	20	1112	Adults	1981- 1982	disease	case control	NJ
Schwartz & Goldstein 1990 [10]	39	633	Adults	1988	disease	case control	NJ
Schwartz et al. 1991 [11]	22	213	Adults	1989	bite	case control	NJ
Orloski et al. 1998 [12]	51	51	All	1993	disease	case control	NJ
Wallace et al. 1998 [13]	19	53	Adults	1995	disease	case control	NY
Townsend et al. 2003 [14]	88	492	All	2001- 2002	disease	case control	NY
Smith et al. 1988 [15]	27	362	Adults	1986	disease	longitudinal	NY
Vaughn & Mesnick 2011 [16]	9	7	Adults	2008	bite	non- controlled trial	NC
Smith et al. 2001 [17]	294	449	All	1998	disease	case control	PA
Connally et al. 2006 [18]	47	47	All	2002- 2003	disease	case control	RI
Finch et al. 2014 [19]	86	809	not specified	2005- 2011	disease	longitudinal	RI
Belongia et al. 2001 [20]	17	45	All	1997- 1998	disease	case control	WI

References

- 1. Aenishaenslin, C., et al., *Exposure and preventive behaviours toward ticks and Lyme disease in Canada: Results from a first national survey.* Ticks Tick Borne Diseases, 2017. **8**(1): p. 112-118.
- 2. Connally, N.P., et al., *Peridomestic Lyme disease prevention: results of a populationbased case-control study.* Am J Prev Med, 2009. **37**(3): p. 201-6.
- 3. Mead, P., et al., *Risk factors for tick exposure in suburban settings in the Northeastern United States.* Ticks and Tick-Borne Diseases, 2018. **9**(2): p. 319-324.
- 4. Vazquez, M., et al., *Effectiveness of Personal Protective Measures to Prevent Lyme Disease*. Emerging Infectious Diseases, 2008. **14**(2): p. 210-216.
- 5. Hinckley, A.F., et al., *Effectiveness of Residential Acaricides to Prevent Lyme and Other Tick-borne Diseases in Humans.* J Infect Dis, 2016. **214**(2): p. 182-8.
- 6. Klein, J.D., S.C. Eppes, and P. Hunt, *Environmental and Life-style Risk Factors for Lyme Disease in Children*. Clinical Pediatrics, 1996. **35**(7): p. 359-363.
- 7. Richards, S.L., J.A.G. Balanay, and J.W. Harris, *Effectiveness of permethrin-treated clothing to prevent tick exposure in foresters in the central Appalachian region of the USA*. International Journal of Environmental Health Research, 2015. **25**(4): p. 453-462.
- 8. Glass, G.E., et al., *Environmental Risk Factors for Lyme Disease Identified with Geographic Information Systems*. American Journal of Public Health, 1994.
- 9. Bowen, G.S., et al., *A Focus of Lyme diseases in Monmouth County, New Jersey.* American Journal of Epidemiology, 1984. **120**(3): p. 387-394.
- 10. Schwartz, B.S. and M.D. Goldstein, *Lyme disease in outdoor workers: risk factors, preventive measures, and tick removal methods.* American Journal of Epidemiology, 1990. **131**(5): p. 877-885.
- Schwartz, B.S., et al., *Antitick saliva antibody a biologic marker of tick exposure that is a risk factor for Lyme disease seropositivity*. American Journal of Epidemiology, 1991.
 134(1): p. 86-95.
- 12. Orloski, K.A., et al., *Emergence of Lyme disease in Hunterdon County, New Jersey,* 1993: a case-control study of risk factors and evaluation of reporting patterns. American journal of epidemiology, 1998. **147**(4): p. 391-7.
- 13. Wallace, B.J., et al., *Human granulocytic ehrlichiosis in New York*. Archives of Internal Medicine, 1998. **158**(7): p. 769-773.
- 14. Townsend, A.K., R.S. Ostfeld, and K.H. Geher, *The effects of bird feeders on Lyme disease prevalence and density of Ixodes scapularis (Acari: Ixodidae) in a residential area of Dutchess County, New York.* Journal of Medical Entomology, 2003. **40**(4): p. 540-546.
- 15. Smith, P.F., et al., *Occupational risk of Lyme-disease in endemic areas of New York state.* Annals of the New York Academy of Sciences, 1988. **539**: p. 289-301.
- 16. Vaughn, M.F. and S.R. Meshnick, *Pilot Study Assessing the Effectiveness of Long-Lasting Permethrin-Impregnated Clothing for the Prevention of Tick Bites.* Vector-Borne and Zoonotic Diseases, 2011. **11**(7): p. 869-875.
- 17. Smith, G., et al., *Risk factors for Lyme disease in Chester County, Pennsylvania*. Public Health Reports, 2001. **116**(SUPPL. 1): p. 146-156.

- 18. Connally, N.P., H.S. Ginsberg, and T.N. Mather, *Assessing peridomestic entomological factors as predictors for Lyme disease*. Journal of Vector Ecology, 2006. **31**(2): p. 364-370.
- 19. Finch, C., et al., *Integrated assessment of behavioral and environmental risk factors for Lyme disease infection on Block Island, Rhode Island.* PLoS One, 2014. **9**(1): p. e84758.
- 20. Belongia, E.A., et al., *Tickborne Infections as a Cause of Nonspecific Febrile Illness in Wisconsin.* Clinical Infectious Diseases, 2001. **54449**.