


RESEARCH

Additional file 1: Supplementary figures

Andrei S Morgan^{1*}, Neil Marlow¹, Elizabeth S Draper³, Zarko Alfirević⁴, Enid M Hennessy⁵ and Kate Costeloe²


*Correspondence: andrei.morgan@ucl.ac.uk
¹Institute for Womens' Health, UCL, 74 Huntley Street, WC1E 6AU, London, UK
 Full list of author information is available at the end of the article

Supplementary figure 1 — Inclusion numbers for the obstetric antecedents analysis




Flow diagram showing inclusion and exclusion numbers for participant mother-baby pairs from the EPICure 2 cohort of babies born between 22 and 26 completed weeks gestation in England in 2006. GA: gestational age.

Supplementary figure 2 — Counselling factors assessed in the obstetric antecedents analysis




Flow diagram showing detail about counselling factors measured in the EPICure 2 cohort of babies born between 22 and 26 completed weeks gestation in England in 2006.

Supplementary figure 3 — Clinical factors assessed in the obstetric antecedents analysis



Flow diagram showing detail about clinical factors measured in the EPICure 2 cohort of babies born between 22 and 26 completed weeks gestation in England in 2006.

Supplementary figure 4 — Delivery information available for the obstetric antecedents analysis



Flow diagram showing detail about delivery measured in the EPICure 2 cohort of babies born between 22 and 26 completed weeks gestation in England in 2006. Not all numbers add to 1,722 due to missing data in some of the variables.

Author details

¹Institute for Womens' Health, UCL, 74 Huntley Street, WC1E 6AU, London, UK. ² Homerton Hospital, London, UK. ³ University of Leicester, Leicester, UK. ⁴ University of Liverpool, Liverpool, UK. ⁵ The Wolfson Institute, Queen Mary University of London, London, UK.

References