

Every Newborn BIRTH multi-country validation study: informing measurement of coverage and quality of maternal and newborn care

Birthweight: EN-BIRTH multi-country validation study

Additional File 15: Adjusted LBW prevalence in exit surveys and routine registers, EN-BIRTH study

		Bangladesh		Nepal	Tanzania	
		Azimpur Tertiary	Kushtia District	Pokhara Regional	Temeke Regional	Muhimbili National
Survey	Unadjusted LBW %	19.8 (18.3, 21.5)	18.1 (16.5, 19.8)	11.1 (10.3, 11.8)	6.7 (6.0, 7.5)	22.0 (20.5, 23.7)
	Adjusted LBW %	21.8 (20.2, 23.5)	21.4 (19.7, 23.2)	14.0 (13.1, 14.8)	9.1 (8.3, 10.0)	23.3 (21.7, 25.0)
Register	Unadjusted LBW %	12.3 (11.0, 13.8)	21.1 (19.2, 23.0)	12.8 (12.0, 13.6)	7.5 (6.9, 8.2)	28.1 (26.6, 29.6)
	Adjusted LBW %	13.5 (12.2, 15.0)	23.9 (22.0, 25.9)	15.5 (14.6, 16.4)	9.6 (9.0, 10.4)	29.2 (27.7, 30.7)

Adjusted LBW prevalence was calculated after re-allocating 25% of 2500g babies to be LBW. Survey n=18,116, register n=20,789.