

**Additional Table 3 Protein levels in PDAC compared to non-neoplastic ductal epithelia**

Marker	High IHC score	PDAC (%)	Duct (%)	p-value
Plasma membrane				
<b>EGFR</b>	$\geq 2$	<b>69</b>	<b>36</b>	<b>0.02</b>
<b>MET</b>	$\geq 1$	<b>65</b>	<b>0</b>	<b>&lt;0.0001</b>
Cytoplasm				
<b>ADAM9</b>	$\geq 3$	<b>88</b>	<b>8</b>	<b>&lt;0.001</b>
<b>SRC</b>	$\geq 2$	<b>58</b>	<b>4</b>	<b>&lt;0.0001</b>
<b><math>\beta</math>CAT</b>	$\geq 2$	<b>38</b>	<b>4</b>	<b>0.005</b>
<b>PKB<math>\beta</math></b>	$\geq 2$	<b>65</b>	<b>16</b>	<b>0.0005</b>
SMAD4	$\geq 2$	38	24	0.4
<b>PTEN</b>	$\geq 1$	<b>42</b>	<b>80</b>	<b>0.01</b>
Nucleus				
<b>SMAD4</b>	$\geq 2$	<b>35</b>	<b>84</b>	<b>0.0005</b>
PTEN	$\geq 1$	50	68	0.2
PKB $\beta$	$\geq 1$	15	4	0.3
<b><math>\beta</math>CAT</b>	$\geq 1$	<b>31</b>	<b>0</b>	<b>0.004</b>

Listed are percentages of specimens with high levels of proteins based on a total of 26 PDAC and 25 non-neoplastic specimens. Highlighted are significant differences in proportions of PDAC compared to ducts in low and high protein expression levels using Fisher's exact test ( $p \leq 0.05$ ).