

Table 1

Treatment	Time intervall after cell injection [min]			
	0 - 10	11 - 20	21 - 30	31 - 40
adherent cells / 20 microscopic fields [mean ±SD]				
T 84				
control IgG [n=6]	16.7 ±9.0	15.8 ±5.1	14.1 ±5.6	13.0 ±1.9
HT-29 LMM				
control IgG [n=9]	13.5 ±5.1	13.5 ±4.8	11.1 ±2.7	13.5 ±2.3
anti integrin-β1[n=7]	11.4 ±4.7	11.8 ±4.4	12.9 ±2.7	12.3 ±3.8
anti integrin-β4 [n=10]	13.4 ±7.6	10.7 ±4.0	10.4 ±3.2	11.0 ±4.5
anti integrin-αv [n=8]	10.6 ±5.1	10.1 ±4.1	9.5 ±2.5	11.7 ±2.7
vehicle [n=12]	12.8 ±4.7	13.2 ±6.7	10.9 ±3.9	15.0 ±5.9
neuraminidase type V [n=8]	7.4 ±3.0	6.6 ±2.3*	6.5 ±2.6*	7.9 ±2.9**
fucoidane (animals) [n=7]	11.6 ±6.9	7.4 ±3.1*	7.2 ±3.3*	5.6 ±2.5**
control medium[n=5]	10.9 ±5.1	18.3 ±10.1	16.4 ±3.9	14.8 ±6.1
anti galectin-3 [n=5]	18.8 ±6.7	13.3 ±2.4	13.1 ±2.9	12.3 ±3.2
anti TF [n=4]	11.5 ±14.7	7.9 ±4.7*	8.6 ±3.7**	7.2 ±4.0*
anti TF + anti galectin-3 [n=5]	11.6 ±1.0	7.4 ±2.8**	7.3 ±2.3**	6.5 ±1.4**

ANOVA and post hoc (LSD):

* p<0.05
 ** p<0.001