

**Table S2. Smoking Opinions and Care by Smoking Status**

Patients' opinions regarding the relationship of smoking and MPN	Patients responding "Yes" (n, %)		
	Current smokers (N=20)	Former smokers (N=161)	Total (N=181)
Smoking Increases Risk of Developing MPN	7 (35.0%)	59 (38.3%)	66 (37.9%)
Chewing Tobacco Increases Risk of Developing MPN	8 (40.0%)	55 (35.5%)	63 (36.0%)
Vaping Increases Risk of Developing MPN	6 (33.3%)	53 (34.6%)	59 (34.5%)
Smoking Increases Risk of Developing Blood Clot	18 (90.0%)	128 (81.5%)	146 (82.5%)
Vaping Increases Risk of Developing Blood Clot	12 (60.0%)	86 (56.2%)	98 (56.6%)
Vaping Safer Than Smoking for MPN Patients	3 (15.8%)	54 (35.5%)	57 (33.3%)
<b>Smoking Cessation Guidance</b>			
Physician Discussed Tobacco Use with You	17 (85.0%)	59 (37.8%)	76 (43.2%)
<b>Medications Attempted to Aid Cessation</b>			
Varenicline	3 (15.0%)	4 (2.5%)	7 (3.9%)
Bupropion	4 (20.0%)	4 (2.5%)	8 (4.4%)
Nicotine Replacements	10 (50.0%)	25 (15.5%)	35 (19.3%)

MPN: Philadelphia-negative chronic myeloproliferative neoplasms