

Additional file 6 Change from baseline to Week 8 in VAS score for sleep disturbance (mITT population)

	Mirogabalin add-on group		Conventional treatment group		Intergroup difference^a
	(N = 63)		(N = 65)		
	Mean ± SD	Change from baseline	Mean ± SD	Change from baseline	
Baseline	46.4 ± 29.6		46.8 ± 31.6		
Day 1	41.1 ± 29.6	-5.3 ± 23.9 <i>P</i> = 0.086 ^b	31.6 ± 24.7	-14.6 ± 33.0 <i>P</i> < 0.001 ^b	<i>P</i> = 0.074
Week 2	16.3 ± 19.7	-31.9 ± 28.3 <i>P</i> < 0.001 ^b	21.8 ± 24.9	-24.0 ± 37.3 <i>P</i> < 0.001 ^b	<i>P</i> = 0.211
Week 4	8.8 ± 13.1	-39.9 ± 28.9 <i>P</i> < 0.001 ^b	15.4 ± 19.8	-30.2 ± 28.3 <i>P</i> < 0.001 ^b	<i>P</i> = 0.081
Week 8	5.6 ± 10.0	-41.4 ± 28.4 <i>P</i> < 0.001 ^b	8.0 ± 12.0	-37.1 ± 32.3 <i>P</i> < 0.001 ^b	<i>P</i> = 0.474

^a vs. the conventional treatment group by *t*-test.

^b vs. baseline by paired *t*-test.

mITT, modified intention-to-treat; SD, standard deviation; VAS, Visual Analogue Scale.