



Haemoglobin < 8 g/dL




Study	Diagnostic OR (95% CI)
Getaneh 2000	7.68 (3.29 - 17.96)
Kalter 1997	7.45 (4.30 - 12.88)
Luby 1995	2.67 (2.00 - 3.58)
Muhe 2000	22.52 (17.19 - 29.50)
Zucker 1997a	8.25 (6.47 - 10.53)
Zucker 1997b	1.04 (0.84 - 1.29)
Zucker 1997c	9.15 (6.89 - 12.14)

Random Effects Model
 Pooled Diagnostic Odds Ratio = 5.94 (2.38 to 14.79)
 Cochran-Q = 377.04; df = 6 (p = 0.0000)



Study	Diagnostic OR (95% CI)
Getaneh 2000	4.02 (1.80 - 9.00)
Kalter 1997	5.32 (3.25 - 8.72)
Luby 1995	5.42 (4.14 - 7.10)
Muhe 2000	34.62 (22.55 - 53.14)
Zucker 1997a	16.24 (12.01 - 21.96)
Zucker 1997b	9.27 (7.14 - 12.03)
Zucker 1997c	13.84 (10.12 - 18.91)

Random Effects Model
 Pooled Diagnostic Odds Ratio = 10.06 (6.16 to 16.41)
 Cochran-Q = 80.37; df = 6 (p = 0.0000)



Study	Diagnostic OR (95% CI)
Getaneh 2000	5.27 (2.34 - 11.84)
Luby 1995	5.24 (4.00 - 6.85)
Muhe 2000	41.40 (26.95 - 63.58)
Zucker 1997a	16.24 (12.01 - 21.96)
Zucker 1997b	14.35 (10.46 - 19.68)

Random Effects Model
 Pooled Diagnostic Odds Ratio = 12.42 (5.97 to 25.85)
 Cochran-Q = 81.22; df = 4 (p = 0.0000)