



Haemoglobin < 5 g/dL




Study	Diagnostic OR (95% CI)
Desai 2002	4.53 (2.43 - 8.44)
Getaneh 2000	11.22 (1.00 - 125.41)
Kalter 1997	30.60 (1.78 - 525.30)
Luby 1995	7.76 (4.36 - 13.82)
Muhe 2000	19.37 (12.60 - 29.79)
Zucker 1997a	33.56 (18.33 - 61.45)
Zucker 1997b	186.63 (64.16 - 542.90)
Zucker 1997c	29.94 (18.99 - 47.20)

Random Effects Model
 Pooled Diagnostic Odds Ratio = 20.62 (10.07 to 42.24)
 Cochran-Q = 55.06; df = 7 (p = 0.0000)



Study	Diagnostic OR (95% CI)
Desai 2002	5.61 (3.02 - 10.43)
Getaneh 2000	5.47 (0.49 - 60.78)
Kalter 1997	7.51 (1.58 - 35.79)
Luby 1995	10.70 (5.00 - 22.91)
Muhe 2000	28.56 (18.38 - 44.38)
Zucker 1997a	90.98 (51.80 - 159.80)
Zucker 1997b	51.40 (29.38 - 89.91)
Zucker 1997c	39.00 (23.81 - 63.87)

Random Effects Model
 Pooled Diagnostic Odds Ratio = 21.62 (10.54 to 44.37)
 Cochran-Q = 62.87; df = 7 (p = 0.0000)



Study	Diagnostic OR (95% CI)
Desai 2002	5.74 (3.09 - 10.67)
Getaneh 2000	7.06 (0.63 - 78.58)
Luby 1995	15.53 (6.58 - 36.63)
Muhe 2000	26.38 (16.99 - 40.94)
Zucker 1997a	107.76 (60.48 - 192.00)
Zucker 1997b	38.40 (24.29 - 60.71)

Random Effects Model
 Pooled Diagnostic Odds Ratio = 22.98 (9.78 to 54.02)
 Cochran-Q = 52.27; df = 5 (p = 0.0000)