LINDSAY (OP2-1)

PHASE B Sim new B

Scenario Script Lindsay		
Case Presentation (to be read to participants):	You are called to attend a vaginal delivery of a 39-week infant because mother is psychiatric (schizophrenia). The mother is 26-year-old, G1 P1. Treatment was Haloperidol Olanzapine, Oxazepam. The pregnancy was uncomplicated, but follow up of the pregnancy began late at 20 weeks. The labour started spontaneously 6 hours ago. The amniotic fluid was clear. The foetal heart rate was normal. The mother wants to leave to go back to the psychiatric hospital. Talk is incoherent The anesthesiology team decided to perform sedation (no opioids). Quick birth after sedation	
Facilitator 1 Midwife	 Bold Point in prebriefing → Make feel it could be an unusual situation. Exit from the room, learners have to prepare what they could need Role in the scenario Come with the newborn, tells she doesn't cry and she is floppy Birthweight is 3000g She has to leave because there is a second delivery in another delivery room If learners stucked, suggests ventilation 	

Pedagogic sheet		
Learning objectives	Objective 1: Initiate resuscitation with correct Phase A	
	Objective 2: Recognition of a newborn requiring PPV and dependant to the PPV	
	Objective 3: Discuss the potential need for intubation	
Stage of the scenario	Initial state: RR 5/min, SpO ₂ 65%, HR 80/min, grunting, limp	
Trigger:		
	Ventilation:	
1-Ventilation	RR 40/min, SpO ₂ 80%, HR 130/min, increase HR after 20 seconds of ventilation	
	No ventilation: leads to bradycardia 80 stop weak cry, RR 0, SpO ₂ 74.	
2- Stop Ventilation	140 ventuation. leads to oracycurdin ou stop weak ery, kit 0, 5p02 /4.	
	Resuming ventilation:	
3 Resuming	RR 55/min, SpO ₂ 90%, HR 150/min at 5 minutes	
ventilation	End of the scenario	
Key points of the	PPV at the right time	
<u>debriefing</u>	 Dependancy to PPV, what to do in this case 	
	Communication between learners	
	o ILCOR Step A	
	0	