

Table 3. Factors associated with minimally adequate treatment for 915 respondents meeting DSM-IV MDE criteria nested in 65 clinics in the Dialogue Project in 2008

	Model with level 1 factors only			Final model		
	Coefficient	OR	95% CI	Coefficient	OR	95% CI
Individual characteristics						
Intercept	-0.72**	0.49	0.31 - 0.78	-1.34***	0.26	0.14 - 0.48
Age						
18-24	-0.43	0.65	0.4 - 1.05	-0.50*	0.61	0.37 - 0.99
25-44 (ref)	-	1.00	-	-	1.00	-
45-64	-0.18	0.83	0.61 - 1.13	-0.20	0.82	0.60 - 1.11
65+	-1.33***	0.27	0.14 - 0.52	-1.45***	0.23	0.12 - 0.47
Sex						
Male	-0.06	0.94	0.68 - 1.3	-0.13	0.88	0.63 - 1.22
Female (ref)	-	1.00	-	-	1.00	-
Has a family physician						
Yes	0.47*	1.61	1.1 - 2.36	0.52**	1.68	1.14 - 2.48
No (ref)	-	1.00	-	-	1.00	-
Has a supplementary insurance coverage						
Yes	0.55***	1.73	1.3 - 2.3	0.53***	1.70	1.27 - 2.27
No (ref)	-	1.00	-	-	1.00	-
At least one comorbid Anxiety Disorder						
Yes	0.58***	1.8	1.35 - 2.39	0.58***	1.79	1.35 - 2.39
No (ref)	-	1.00	-	-	1.00	-
Severity of depressive symptoms (HADS depression sub-scale)	0.07***	1.07	1.04 - 1.11	0.07***	1.07	1.04 - 1.11
Clinics characteristics						
Psychotherapy available on-site						
Yes				0.38*	1.46	1.06 - 2.02
No (ref)				-	1.00	-
Number of GP using treatment algorithms with individuals suffering from anxiety or depressive disorders						
All/Most				0.40*	1.49	1.07 - 2.08
None/Some (ref)				-	1.00	-
Inadequate mode of remuneration to offer an “optimal level” of care for patients suffering from anxiety or depressive disorders						
Not at all / Slightly				0.46*	1.58	1.07 - 2.35
Fairly				0.29	1.33	0.91 - 1.95
Highly (ref)				-	1.00	-

* $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$

