

Table 2: Participant characteristics, diagnosis, and tobacco consumption patterns (n=12)

Name ^a	SC ^b	Age	Cigarettes per day at time of interview	Nicotine dependence ^c	ADHD subtype	ICD 10 comorbidity	Use of other substances	Current medication	Previous phases of nicotine abstinence
Mr. A.	yes	53	2 - 3	very low	combined	-	-	stimulants	1 month (1993)
Mr. B.	yes	42	3 – 5 / week	very low	inattentive	F33.4	-	stimulants, antidepressants	7 years (starting 1996)
Mrs. C.	yes	25	20	low	combined	F12.24	cannabis	stimulants, antidepressants	15 months (2002)
Mrs. D.	yes	46	20	high	combined	F10.25 F11.20 F14.20	alcohol	stimulants, antidepressants	6 months (1993)
Mrs. E.	yes	27	20	very high	combined	-	-	stimulants	4 – 5 years (2000)
Mrs. F.	yes	47	23	very high	combined	F43.21	-	stimulants, antidepressants	3 weeks (1994)
Mrs. G.	yes	42	30	moderate	combined	F43.21	-	Antidepressants	One year (2001)
Mrs. H.	no	26	35	high	combined	F43.21	-	stimulants, antidepressants	18 months (2003)
Mr. I.	no	47	35	high	hyperactive/impulsive	F12.24 F60.2	cannabis	stimulants	1 month (2003)
Mrs. J.	yes	44	30	very high	combined	F11.202 F41.2	alcohol	stimulants	6 months (1984)
Mr. K.	no	46	33	very high	combined	F10.24 F33.4	alcohol	stimulants	1 month (1992)
Mr. L.	no	32	35	very high	combined	F10.25 F14.20	alcohol	stimulants	none

^a pseudonym

^b participated in smoking cessation treatment

^c nicotine dependence according to Fagerström