

Appendix:

Table e1. Antidepressants taken by respondents

% of patients on...		Total	US	Canada	UK	Germany	France	Spain
Minimal dose (mg)		N=2,096	n=597	n=301	n=300	n=299	n=299	n=300
SSRI		76.7%	69.8%	78.1%	70.3%	72.2%	87.3%	89.0%
	citalopram 20							
	escitalopram 10							
	fluoxetine 20							
	fluvoxamine 100							
	paroxetine 20							
	paroxetine CR 37.5							
	sertraline 100							
SNRI		33.8%	22.4%	41.5%	21.0%	39.1%	40.1%	49.7%
	desvenlafaxine 50							
	duloxetine 60							
	levomilnacipran 40							
	milnacipran 100							
	venlafaxine 75							
MAOI		6.8%	0.2%	9.3%	2.7%	14.4%	14.0%	0.0%
	phenelzine 60							
	tranylcypromine 30							
TCA		18.8%	5.2%	17.3%	21.7%	26.8%	25.8%	29.3%
	amitriptyline 75							
	clomipramine 100							
	desipramine 100							
	imipramine 75							
	nortriptyline 75							
OTHER ADTs		24.2%	23.8%	25.6%	20.3%	31.1%	18.1%	27.0%
	agomelatine 25							
	bupropion 200							
	mirtazapine 30							
	trazodone 150							
	vilazodone 20							
	vortioxetine 20							

Figure e1. Symptom frequency (a) individual PHQ-9 items (b) symptoms not assessed by PHQ-9 (n=2,096)

