

**Table S1***Demographic and clinical variables of participants (N=192)*

	<i>N</i> <sup>a</sup> (%) <sup>b</sup>
<b>Gender</b>	
Male	134 (69.8)
<b>Age group</b>	
18-24	24 (12.5)
25-34	79 (41.1)
35-44	49 (25.5)
45-54	28 (14.6)
55+	12 (6.3)
<b>Marital status</b>	
Partnered	117 (60.9)
<b>Mode of Arrival</b>	
Irregular maritime arrival	104 (54.2)
<b>Continent of Origin (UN geoscheme)</b>	
Africa	
East Africa	17 (9.1)
North Africa	10 (5.3)
West Africa	7 (3.7)
Asia	
South Asia	97 (51.9)
South-East Asia	39 (20.9)
West Asia	8 (4.3)
Other	9 (4.7)
<b>Interpreter required</b>	
Yes (screening)	118 (61.5)
Yes (interview)	125 (65.1)
<b>Pre-migration camp/detention</b>	
Yes	28 (14.8)
<b>Post-migration Immigration Detention</b>	
Yes	105 (54.7)
<b>Years in Australia</b>	
< 1 year	36 (19.0)
1-2 years	34 (18.0)
2-3 years	79 (41.8)
> 3 years	40 (21.2)
<b>Residency status</b>	
Temporary Visa (Asylum-seeker)	187 (97.4)
Permanent Residency (Refugee)	5 (2.6)
<b>Mental health diagnosis (Australia)</b>	
Yes	8 (4.4)
<b>Post-traumatic stress disorder (PTSD)<sup>c</sup></b>	

Yes	38 (19.9)
<b>Major depressive disorder (MDD)</b>	
Yes	57 (29.7)
<b>Either PTSD or MDD</b>	
Yes	62 (32.3)

---

<sup>a</sup> Total *ns* may be less than 192 due to missing data

<sup>b</sup> Refers to valid percentage, excluding missing data.

<sup>c</sup> *n* = 191.