

Supplementary table 1: Summary of between group differences of onset of basic symptoms compared to onset of attenuated psychotic symptoms

	<u>Paired t-test</u>						<u>Robust regression</u>					
	<b>N</b>	<b>df</b>	<b>t</b>	<b>MD</b>	<b>p</b>	<b>95% CI</b>	<b>Cohen's d</b>	<b>B</b>	<b>SE</b>	<b>t-value</b>	<b>F</b>	<b>p</b>
Male only	19	18	-0.01	-0.004	1.00	-0.69, 0.68	0.00	-0.055	0.188	-0.293	0.09	0.770
Female only	45	44	0.04	0.007	1.00	-0.36, 0.37	-0.01	0.094	0.093	1.021	1.02	0.319
< 18 years	41	40	0.1	0.02	0.80	-0.4, 0.44	0.15	-0.015	0.136	-0.113	0.01	0.911
> 18 years	23	22	-0.1	-0.026	0.90	-0.53, 0.48	-0.10	-0.016x	0.085	-0.193	0.04	0.847
No 3rd level education	9	7	0.4	0.18	0.70	-0.78, 1.14	-0.22	-0.037	0.195	-0.191	0.04	0.855
3rd level education	55	54	-0.1	-0.021	0.90	-0.37, 0.33	0.02	0.068	0.094	0.722	0.5	0.472