

**Item score distributions of the PANSS-8 (N = 1744)**

<b>Item</b>	<b>Mean (SD<sup>a</sup>)</b>	<b>Median (range)</b>	<b>Floor<sup>b</sup></b>	<b>Ceiling<sup>c</sup></b>
<b>P1 Delusions</b> <i>(positive symptom, psychoticism)</i>	2.37 (1.65)	2 (1–7)	47%	1%
<b>G9 Unusual thought content</b> <i>(general symptom, psychoticism)</i>	2.00 (1.38)	1 (1–7)	56%	1%
<b>P3 Hallucinatory behaviour</b> <i>(positive symptom, psychoticism)</i>	1.99 (1.50)	1 (1–7)	62%	0%
<b>P2 Conceptual disorganization</b> <i>(positive symptom, disorganization)</i>	1.95 (1.33)	1 (1–7)	56%	0%
<b>G5 Mannerisms/Posturing</b> <i>(general symptom, disorganization)</i>	1.59 (1.03)	1 (1–7)	68%	0%
<b>N1 Blunted affect</b> <i>(negative symptom)</i>	2.07 (1.29)	2 (1–7)	48%	0%
<b>N4 Social withdrawal</b> <i>(negative symptom)</i>	2.38 (1.54)	2 (1–7)	42%	1%
<b>N6 Lack of spontaneity</b> <i>(negative symptom)</i>	2.04 (1.38)	1 (1–7)	53%	0%
<b>Total score</b>	16.39 (7.76)	15 (8–51)	16%	0%

<sup>b</sup>% of lowest possible score

<sup>c</sup>% of highest possible score

The sub-classification of items according to two different models (positive/negative/general symptoms or psychoticism/disorganization/negative symptoms) is indicated in parentheses.