

Table 5: Use of condoms with three types of sex partners in the previous year according to HIV serostatus

Use of condoms with	HIV (+)	HIV (-)	Total	p-value* (χ^2 test)
Wife or girlfriend (valid response: n=118)				
Every time	2 (5.4)	5 (6.2)	7 (5.9)	0.91
Almost every time	1 (2.7)	3 (3.7)	4 (3.4)	
About half	2 (5.4)	3 (3.7)	5 (4.2)	
Sometimes	3 (8.1)	11 (13.6)	14 (11.9)	
Never	29 (78.4)	59 (72.8)	88 (74.6)	
Casual partners (valid response: n=19)				
Every time	2 (33.3)	4 (30.8)	6 (31.6)	0.85
Almost every time	0 (0.0)	1 (7.7)	1 (5.3)	
About half	0 (0.0)	0 (0.0)	0 (0.0)	
Sometimes	1 (16.7)	1 (7.7)	2 (10.5)	
Never	3 (50.0)	7 (53.8)	10 (52.6)	
FSWs (valid response: n=51)				
Every time	7 (46.7)	16 (44.4)	23 (45.1)	0.28
Almost every time	1 (6.7)	10 (27.8)	11 (21.6)	
About half	1 (6.7)	0 (0.0)	1 (2.0)	
Sometimes	3 (20.0)	5 (13.9)	8 (15.7)	
Never	3 (20.0)	5 (13.9)	8 (15.7)	

n indicates number; IDU, injection drug user

*: Differences in proportions between HIV (+) IDUs and HIV (-) IDUs were tested by χ^2 tests.