




	Prevalence Year 2000	Prevalence Year 2008
Urban	7.5 (6.6-8.4)	8.4 (7.0-9.8)
Rural	5.4 (4.2-6.6)	9.7 (7.6-11.8)
	<sup>1</sup> p=0.01	<sup>1</sup> p=0.33




	Prevalence Year 2000	Prevalence Year 2008
Urban	11.9 (10.8-13.1)	17.1 (15.1-19.1)
Rural	9.7 (8.1-11.3)	18.4 (15.6-21.2)
	<sup>1</sup> p=0.03	<sup>1</sup> p=0.49




	Prevalence Year 2000	Prevalence Year 2008
Urban	3.6 (3.0-4.2)	5.2 (4.0-6.4)
Rural	3.7 (2.7-4.7)	4.0 (2.6-5.4)
	<sup>1</sup> p=0.84	<sup>1</sup> p=0.22



	Prevalence Year 2000	Prevalence Year 2008
Urban	6.8 (6.0-7.6)	14.8 (12.9-16.7)
Rural	6.9 (5.5-8.3)	11.9 (9.6-14.2)
	<sup>1</sup> p=0.83	<sup>1</sup> p=0.08



	Prevalence Year 2000	Prevalence Year 2008
Urban	2.3 (1.8-2.8)	2.3 (1.5-3.1)
Rural	1.5 (0.8-2.2)	3.0 (1.8-4.2)
	<sup>1</sup> p=0.15	<sup>1</sup> p=0.34



	Prevalence Year 2000	Prevalence Year 2008
Urban	2.9 (2.4-3.4)	4.8 (3.7-5.9)
Rural	1.9 (1.2-2.6)	6.0 (4.3-7.7)
	<sup>1</sup> p=0.08	<sup>1</sup> p=0.24

<sup>1</sup>p value of chi-square test comparing the prevalence of asthma and allergy outcomes between urban and rural areas within each survey year.