Supplemental Table 1: Distribution of socio-demographic, sexual behaviour and sex-work related characteristics, clients of female sex workers, Karnataka, South India $(N=2,757)^a$

	Characteristic	%
Socio-demographic	Age	
	18-24	29.7
	25-29	24.6
	30-39	28.7
	40+	17.0
	Mean	30.4
	Marital status	
	Married	62.1
	Separated	2.6
	Never married	35.3
	Occupation	25.0
	Transport	25.9
	Service	13.8
	Business Other (including Labourer)	11.6
	Other (including Labourer) Can read and write	48.7
	Can read and write No	25.3
	Yes	23.3 74.7
	Districts	/4./
	Belgaum	17.3
	Bagalkot	17.1
	Bellary	14.2
	Shimoga	15.5
	Bangalore	30.6
	Mysore	5.4
Sexual Behaviour	Age at 1 st sex	
	<18	15.4
	18-20	52.9
	21-24	21.1
	25+	10.6
	Mean	20.0
	Have an intimate partner	66.2
	Condom use in last sex with intimate partner	66.2
	Condoni use in fast sex with intimate partier	10.3
	Never use condoms with intimate partner	10.5
	rever use condoms with miniate partier	86.5
	Anal sex with intimate partners	00.0
		4.6
	Anal sex with MSM/hijra (6 months)	
	J (/	5.5

Sex-Work Related	Age at 1 st paid sex	
	<18	7.4
	18-20	47.2
	21-24	26.7
	25+	18.7
	Mean	21.5
	Duration of paid sex	
	1 year or less	14.3
	2-4	25.3
	5-9	22.4
	10+	38.0
	Mean	8.9
	Place of solicitation, FSWs	
	Public places	36.0
	Brothel	9.8
	Home	47.8
	Lodge	6.4
	Number of FSWs (Total: 6 months)	
	1	17.0
	2-3	40.7
	4-5	22.2
	6-9	12.6
	10+	7.4
	Mean	4.3
	Number of Occasional FSWs (6 months)	
	0	14.3
	1	12.9
	2-4	51.8
	5+	21.1
	Mean No. 1 Favy (6 11)	3.4
	Number of Regular FSWs (6 months)	57 .0
	0	57.8
	1	28.5
	2-4	10.4
	5+	3.4
	Mean Condenses in less consistent ESW	0.9
	Condom use in last sex, occasional FSW	66.0
	Never use condems, conscional ESW	66.9
	Never use condoms, occasional FSW	26.2
	Condom use in last say regular ESW	26.3
	Condom use in last sex, regular FSW	50.0
	Navar usa condoms regular ESW	59.9
	Never use condoms, regular FSW	32.4
	Ever asked for anal intercourse, FSW	34.4
	Evel asked for aliai illefedurse, I's w	11.2
		11.2

	Ever had anal sex, FSW	
	Condom use in last anal sex, FSW	7.8
	Condoni use in last and sex, 15 w	57.1
Pathogen Prevalence	CT	
J	NC	1.8
	NG	0.5
	CT/NG	2.2
	Active syphilis	
	HIV	3.6
		5.6
	HSV-2	28.4

^aMSM: Men who have sex with men; FSW: Female sex workers; CT: Chlamydia; NG: Gonorrhea; HIV: Human immunodeficiency virus; HSV-2: Herpes simplex virus, type 2