

## Tables

**Table 1: Relationship between risk factors of water supply status among PLWHA on Home Based Care clients of Gondar town, 2009. (n= 294)**

Characteristics		General Water supply			
		status		Crude OR (95% CI)	Adjusted OR (95% CI)
		Unimproved	Improved		
Educational status	Illiterate(can't Read and write)	58	71	1.0	1.0
	Read and write	27	14	2.36(1.13,4.91)*	4.20(0.92, 19.02)
	Elementary	17	34	0.61(0.31,1.20)	0.74(0.19, 2.78)
	Secondary	19	39	0.59(0.31,1.14)	0.50(0.17, 1.46)
	Above grade12	5	10	0.61(0.19,1.89)	0.16(0.01, 2.15)
Income quintile	Lowest( poorest)	15	38	0.47(0.20,1.081)	0.67(0.137, 3.36)
	Second (poor)	11	30	0.43(0.17,1.706)	0.39(0.057, 2.77)
	Third (middle)	52	48	1.29(0.64,2.59)	3.55(0.96, 13.13)
	Fourth(High)	27	27	1.19(0.54,2.61)	2.99(0.72, 12.42)
	Fifth (Highest)	21	25	1.0	1.0
Discrimination of family member in water source	Yes	9	9	1.35 (0.52, 3.52)	0.43 (0.06, 2.84)
	No	117	159	1.0	1.0
Discrimination of PLWHA in water source	Yes	13	22	0.76(.36, 1.58)	1.61 (0.44, 5.86)
	No	113	146	1.0	1.0
Forced to go far distance to fetch water	Yes	9	4	3.91(1.13, 13.47)*	3.84 (0.41, 35.27)
	No	42	73	1.0	1.0

do you need	yes	1	11	0.11(0.01, 0.89)*	0.13(0.01, 1.44)
help with	No	125	157	1.0	1.0
walking					
diarrhea for		15	8	2.70(1.10, 6.59)*	6.13(1.23, 30.57)*
the past 24	Yes				
hour	No	111	160	1.0	1.0

---

\* Statistically significant association p<0.05