Table S2. Caregivers' experiences with external sources^a

	Positive Outcome: Child Recovered			Negative Outcome: Child Survived but with Def			Negative Outcor	
$Cases \to$	CS 06	CS 07	CS 08	CS 01	CS 02	CS 03	CS 04	CS 05
1st Action	Nearest PHC Accessed:	Nearest PHC Accessed:	Nearest PHC Accessed: Yes	Nearest PHC Accessed:	Nearest PHC Accessed: Yes	Nearest PHC Accessed:	Nearest PHC Accessed: Yes	Nearest PHC Accessed: Yes
	Reason not using PHC: Stock-outs, told to go to drug shop, health professionals are rude, & cost of transport Used: Drug shop ~ 1 mile	Reason not using PHC: Lack of staff, stock-outs, & cost of transport Used: Drug shop ½ mile	Used: Muhuyu HC II	Reason not using closest PHC: stock-outs Used: Kalabangha HC III ~ 5 miles	Used: Busaba HC III ~ 3 miles	Reason not using PHC: verbal abuse from health professionals, no injections given, lack of examination & stock-outs Used: Drug Shop ~ 3 miles	Used: Nabiganda HC II	Used: Nakwasi HC III
	Setting: Drug shop Experience: Took child ~ 12 hours from start of symptoms No blood test done No information about illness given Vendor suggested antimalarial syrup, but refused as had no money Caregiver requested Aspirin® for 3 days. No money to buy more drugs Cost was 100UGX, willing to pay up to 500UGX Continued to be sick but no additional medicines given. Gradually resolved	Setting: Drug shop Experience: Took child ~ 9 hours from start of symptoms Blood test done but not told of results No information about illness given Bought medicines Cost was 2000UGX Improved gradually	Setting: Muhuyu HC II Experience: Took child after 1 day from start of symptoms Had to wait 4 hours before being seen Was examined No blood test done No information about illness given Received free medicines Cured in a few days	Setting: Kalabangha HC III Experience: Took child ~ 12 hours from start of symptoms Was examined No blood test done Told was anemic, had high fever No medicines given Referred to hospital in neighboring district of Mbale because had no blood to transfuse Subsequent action taken	Setting: Busaba HC III Experience: Took child ~ 6 hours from start of symptoms Waited 30 minutes before health professionals arrived Blood test done Told had malaria & convulsions from fever Received free medicines & told to buy more from drug shop Not improved, convulsions worsened Subsequent action taken	Setting: Drug shop Experience: Took child ~ 17 hours from start of symptoms Provider was busy but was provided immediate care because saw child was in a bad state, told others to wait while she treated the child No blood test done Examined, touched & looked at eyes Told to cover child with blanket, hold him so doesn't fall down, & to return the next day Bought medicines Cost was 13000UGX for day 1; 7500UGX for day 2 Malaria resolved gradually, but stomach remained swollen & child continued to convulse	Setting: Nabiganda HC II Experience: Took child ~ 15 hours from start of symptoms No blood test done Told was anemic Received no medicines Told would not treat because needed to first receive blood transfusion Child referred to hospital in neighboring district of Mbale Subsequent action taken	Setting: Nakwasi HC III Experience: Took child ~ 1 hour from start of symptoms No blood test done No information about illness given Received free medicines & told to buy more from drug shop Did not improve Cost not shared Not improved, body swollen, could not eat Subsequent action taken
2 nd Action				Experience: Health professionals ignored her child even when asked for help Blood test done but not told of results Told to buy water for drip, quinine injections, quinine syrup for 2 weeks and other oral medicines, and then to come back Cost was 15000UGX Malaria resolved, but eye sight affected, could not see, left hand became lame, experiences convulsions regularly	Setting: Drug shop & visited Busaba HC III & Busolwe hospital x 1 year Experience: Gave quinine syrup for 1 year on advice from neighbors, health professional, & drug vendor. But stopped when child stopped talking At Busaba HC III & Busolwe hospital given quinine injections, & bought quinine syrup from drug shop every 2 weeks Cost 2500UGX every 2 weeks At times sent away from Busaba HC III because of stock-outs At Busaba HC III & Busolwe hospital never referred for convulsions Blood test never repeated Malaria resolved, but left unable to speak and brain-damaged		Experience: Took blood test but not told of results Caregiver asked to get blood cross-matched at the laboratory, then to buy blood & transfusion kit Child died before blood could be purchased Health professional gave no information about illness No treatment started during the 3 hours at the hospital, arrived 11 am & died at 1 pm After child died, body cleaned & wrapped, returned to family	Experience: Returned in 2 weeks, then frequently over 4 months. Last visit at drug shop because HC was closed Would improve some, but illness would come back Same treatment at HC as 1st action Free medicines & told to buy more from drug shop No blood test No information about illness given Last visit at drug shop, given tablets for worms child died later that day

^aAbbreviations: Public Health Centre (PHC); Health Centre (HC); Ugandan Shilling (UGX).