Additional file 3 Comparisons between non-participation and participation at Wave2 – Wave4 by predictors at baseline

N = 6,182	Wave 2 (N = 6,154)			Wave $3 (N = 6,110)$			Wave 4 $(N = 6,022)$		
	non-Part.	Part.		non-Part.	Part.		non-Part.	Part.	
	(n=1,886)	(n=4,268)	<i>p</i> -value	(n=2,686)	(n=3,424)	<i>p</i> -value	(n=3,615)	(n=2,407)	<i>p</i> -value
Covariates	%/Mean	%/Mean		%/Mean	%/Mean		%/Mean	%/Mean	
Gender									
Female	53.6%	55.4%	0.188	52.6%	56.4%	0.003	53.7%	56.3%	0.047
Male	46.4%	44.6%		47.4%	43.6%		46.3%	43.7%	
Age (years)	54.99	52.82	< 0.001	54.75	52.46	< 0.001	54.00	52.68	< 0.001
30 ~ 64	75.0%	82.5%	< 0.001	75.3%	84.1%	< 0.001	77.8%	83.9%	< 0.001
<b>≧</b> 65	25.0%	17.5%		24.7%	15.9%		22.2%	16.1%	
Education (years)	9.24	10.44	< 0.001	9.61	10.48	< 0.001	9.88	10.44	< 0.001
No schooling	13.1%	6.4%	< 0.001	11.4%	6.0%	< 0.001	10.2%	5.5%	< 0.001
below Senior HS	41.1%	38.0%		39.7%	38.1%		38.9%	38.5%	
Senior HS	27.7%	32.1%		28.5%	32.7%		28.5%	34.6%	
above Senior HS	18.1%	23.5%		20.5%	23.2%		22.4%	21.5%	
Marital status									
Living with spouse	85.7%	90.4%	< 0.001	86.9%	90.5%	< 0.001	87.4%	91.0%	< 0.001
Single	14.3%	9.6%		13.1%	9.5%		12.6%	9.0%	
Tobacco smoking	21.6%	16.4%	< 0.001	21.1%	15.5%	< 0.001	19.5%	15.6%	< 0.001
(Yes)									
Betel-nuts chewing	4.7%	2.5%	< 0.001	4.2%	2.4%	< 0.001	3.9%	2.1%	< 0.001
(Yes)									
Psychiatric status (by CHQ-12)	3.05	2.95	0.040	2.98	2.99	0.825	2.97	3.00	0.462

Normal	70.4%	75.2%	< 0.001	73.3%	74.0%	0.563	73.4%	74.3%	0.421	
Psychiatric	29.6%	24.8%		26.7%	26.0%		26.6%	25.7%		
disorder*										
# of 15 diseases <sup>†</sup>	1.79	1.54	< 0.001	1.77	1.49	< 0.001	1.72	1.45	< 0.001	
# of 4 Pills taken <sup>‡</sup>	0.09	0.10	0.669	0.10	0.10	0.669	0.10	0.09	0.667	
Regular exercise										
status										
No exercise	43.1%	34.3%	< 0.001	41.6%	33.3%	< 0.001	39.5%	33.0%	< 0.001	
Irregular exercise	37.8%	40.9%		39.3%	40.6%		39.6%	40.7%		
<150 minutes	8.2%	11.2%		8.9%	11.5%		9.0%	12.4%		
150~299 minutes	5.0%	6.1%		4.7%	6.6%		5.3%	6.4%		
$\geq 300 \text{ minutes}$	5.8%	7.5%		5.6%	8.1%		6.6%	7.6%		
PCS score	52.25	53.38	< 0.001	52.41	53.52	< 0.001	52.69	53.56	< 0.001	
MCS score	49.52	49.70	0.492	49.67	49.64	0.920	49.45	50.00	0.024	
Attrition										
Died	21		48			88				
Lost to follow-up	7			24			72			

<sup>\*</sup> CHQ-12 score ≥ 4

<sup>&</sup>lt;sup>†</sup> 15 self-reported and/or diagnosed chronic diseases, including T2DM, hypertension, hyperlipidemia, kidney disease, cardiac disease, stroke, hepatic disease, gout, osteoporosis, asthma, psychiatric disease, nerve-related disease, intestinal disease, Tuberculosis, and metabolic syndrome

<sup>&</sup>lt;sup>‡</sup> 4 pills, including refreshing drugs, sleeping pills, sedative medicine, and painkiller *Abbreviations*: PCS, physical component summary; MCS, mental component summary; T2DM, type 2 Diabetes mellitus; non-Part., non-Participation; Part., Participation