

Supplementary Table 3. Hazard ratio for all-cause death between combinations of categories of hard and light physical activity levels over as time-dependent covariates.

		Hazard ratio [95% Conf, Interval]	
Light physical activity levels			
None		0,97	(0,67, 1,42)
<1 hour per week		1,38	(0,89, 2,13)
1-2 hours per week		1,18	(0,87, 1,62)
>3 hours per week			
Hard physical activity levels			
None		1,56	(1,34, 1,83)
Low		1,16	(0,96, 1,40)
Moderate		1,06	(0,88, 1,27)
Vigorous			
Interaction			
Light physical activity levels	Hard physical activity levels		
<i>None</i>	<i>None</i>	1,57	(1,07, 2,31)
<i>None</i>	<i>Some</i>	1,20	(0,62, 2,30)
<i>None</i>	<i>Moderate</i>	0,88	(0,50, 1,54)
<i><1 hour per week</i>	<i>None</i>	0,85	(0,54, 1,32)
<i><1 hour per week</i>	<i>Some</i>	0,65	(0,40, 1,07)
<i><1 hour per week</i>	<i>Moderate</i>	0,59	(0,33, 1,03)
<i>1-2 hours per week</i>	<i>None</i>	0,87	(0,63, 1,20)
<i>1-2 hours per week</i>	<i>Some</i>	0,73	(0,51, 1,04)
<i>1-2 hours per week</i>	<i>Moderate</i>	0,81	(0,56, 1,17)
<i>Test statistics: LR chi2(17) = 10633.14, p<0.0001</i>			