		Baseline		Immediate follow-up		3 month
		(n=50)		(n=35)		( <b>n=30</b> )
	N	Mean (SD)	n	Change from baseline	n	Change from baseline
				Mean (SD)		Mean (SD)
Risk Perception						
Perceived AR	45	29.3 (22.8)	31	-4.1 (-14.0 to 5.8)	22	-0.52 (-8.45 to 7.41)
Conviction of Perceived AR	48	3.58 (1.7)	32	0.33 (-0.45 to 1.1)	26	0.33 (-0.31 to 0.96)
Perceived RR	50	3.48 (1.2)	35	0.2 (-0.24 to 0.64)	29	0.03 (-0.51 to 0.58)
Conviction of Perceived RR	47	3.86 (1.6)	33	0.20 (-0.41 to 0.82)	24	0.38 (-0.25 to 1.00)
Risk Perception						
Perceived AR	45	29.3 (22.8)	31	-4.1 (-14.0 to 5.8)	22	-0.52 (-8.45 to 7.41)
Conviction of Perceived AR	48	3.58 (1.7)	32	0.33 (-0.45 to 1.1)	26	0.33 (-0.31 to 0.96)
Perceived RR	50	3.48 (1.2)	35	0.2 (-0.24 to 0.64)	29	0.03 (-0.51 to 0.58)
Conviction of Perceived RR	47	3.86 (1.6)	33	0.20 (-0.41 to 0.82)	24	0.38 (-0.25 to 1.00)
Maladaptive Coping Behaviour	48	9.3 (2.0)	35	0.35 (-0.47 to 1.18)	29	0.45 (-0.39 to 1.28)
Cancer Worry	46	4.8 (2.0)			25	0.12 (-0.40 to 0.64)
Anxiety	45	12.1 (2.2)	34	0.75 (-0.14 to 1.64)	25	0.64 (-0.22 to 1.50)
		% (95% CI)		% (95% CI)		% (95% CI)
Absolute risk accuracy	45		31		24	
Accurate (±5%)		17.8 (8.9 to 32.4)		25.8 (12.9 to 44.9)		25.0 (11.0 to 47.4)
Underestimate		4.4 (1.1 to 16.9)		0 (0)		0 (0)
Overestimate		77.8 (62.8 to 87.9)		74.2 (55.1 to 87.1)		75.0 (52.6 to 89.0)
Comparative risk accuracy	50		35		29	
Accurate		36.0 (23.6 to 50.6)		25.7 (13.5 to 43.4)		41.4 (24.3 to 60.8)
Underestimate		48.0 (34.2 to 62.1)		51.4 (34.5 to 68.0)		44.9 (27.2 to 63.9)
Overestimate		16.0 (8.0 to 29.4)		22.9 (11.4 to 40.5)		13.8 (4.9 to 33.0)

## Additional File 8. Psychological outcomes and risk factor awareness within the groups receiving the intervention