

Table 3: Psychometric properties of the newly adapted Stigma scale			
Validity	Correlations		
CES-D (<i>n, r, p-value</i>)	<i>n</i> = 44, <i>r</i> = 0.529, <i>p</i> < .001		
GAD-7 (<i>n, r, p-value</i>)	<i>n</i> = 39, <i>r</i> = 0.371, <i>p</i> = .014		
CF symptom (<i>n, r, p-value</i>)	<i>n</i> = 44, <i>r</i> = 0.479, <i>p</i> = .001		
SF- 12 (<i>n, r, p-value</i>)	<i>n</i> = 44, <i>r</i> = -.454, <i>p</i> = .002		
CF-specific QOL (<i>n, r, p-value</i>)	<i>n</i> = 39, <i>r</i> = -.645, <i>p</i> < .001		
Reliability			
Internal consistency (α)	0.79		
Mean Inter-item correlation (mean, N)	0.30 (44)		
Three month test-retest (<i>r, p-value</i>)	<i>r</i> =0.078; <i>p</i> <.001		
	Baseline	Follow-up	p-value
Mean differences between response items (mean, SD)	16.6 (4.5)	16.9 (5.1)	0.3

Center for Epidemiologic Study-Depression scale (CES-D)

Cystic Fibrosis symptom scale (CF symptom)

General Anxiety Disorder -7 scale (GAD-7)

Short Form (SF)-12

Cystic Fibrosis-specific Quality of Life (CF-specific QOL)