

**Additional file 2.** Ischemic ECG changes in all subjects (n=1625), comparing non-COPD and COPD

		Non-COPD n=991	COPD n= 634	P
Ischemic ECG changes				
	Q-Waves	Any, n (%)	78 (7.9)	54 (8.5)
		Major Q/QS, n (%)	33 (3.5)	26 (4.3)
		Minor Q/QS, n (%)	45 (4.7)	28 (4.6)
	ST-segment depressions	Any, n (%)	69 (7.0)	36 (5.7)
		Major, n (%)	6 (0.6)	4 (0.7)
		Intermediate, n (%)	34 (3.6)	21 (3.4)
		Minor, n (%)	21 (3.1)	11 (1.8)
	T- wave items	Any, n (%)	173 (17.5)	110 (17.4)
		Major, n (%)	3 (0.4)	1 (0.2)
		Intermediate, n (%)	74 (8.3)	49 (8.6)
		Minor, n (%)	96 (10.5)	60 (10.3)
	Bundle branch block	LBBB <sup>1</sup> , n (%)	18 (2.1)	15 (2.8)
Whitehall criteria				
	Ischemic heart disease	Any, n (%)	240 (24.2)	158 (24.9)
		Probable <sup>2</sup> , n (%)	48 (6.0)	40 (7.8)
		Possible <sup>3</sup> , n (%)	192 (20.4)	118 (19.9)

<sup>1</sup>Left bundle branch block. <sup>2</sup>Including Major Q/QS and LBBB. <sup>3</sup>Including Minor Q/QS, any ST-segment depression and any T-wave item