Study or Subgroup	post Mean	post SD	post Total	pre Mean	pre SD	pre Total	Weight	Mean Difference IV, Random, 95% CI
4.2.1 vs wlc								
Subtotal (95% CI)			**0**			**0**		**Not estimable**
Heterogeneity: Not applicable								
Test for overall effect: Not applicable								
4.2.2 vs sham ++								
Basford 2000	41.6	39	23	61.1	67.5	23	3.6%	-19.50 [-51.36, 12.36]
Subtotal (95% CI)			**23**			**23**	**3.6%**	**-19.50 [-51.36, 12.36]**
Heterogeneity: Not applicable								
Test for overall effect: Z = 1.20 (P = 0.23)								
4.2.3 vs sham -								
Lam 2007	30.5	17.7	21	51.4	18.8	21	30.0%	-20.90 [-31.94, -9.86]
Sterigioulas_2007	19.79	9.11	25	38.14	18.46	31	66.4%	-18.35 [-25.76, -10.94]
Subtotal (95% CI)			**46**			**52**	**96.4%**	**-19.14 [-25.30, -12.99]**
Heterogeneity: Tau² = 0.00; Chi² = 0.14, df = 1 (P = 0.71); I² = 0%								
Test for overall effect: Z = 6.09 (P < 0.00001)								
Total (95% CI)			**69**			**75**	**100.0%**	**-19.16 [-25.20, -13.11]**
Heterogeneity: Tau² = 0.00; Chi² = 0.14, df = 2 (P = 0.93); I² = 0%								
Test for overall effect: Z = 6.21 (P < 0.00001)								
Test for subgroup differences: Chi² = 0.00, df = 1 (P = 0.98), I² = 0%