Study or Subgroup	post Mean	post SD	post Total	pre Mean	pre SD	pre Total	Weight	Mean Difference IV, Random, 95% CI
4.3.1 vs wlc								
Subtotal (95% CI)			**0**			**0**		**Not estimable**
Heterogeneity: Not applicable								
Test for overall effect: Not applicable								
4.3.2 vs sham ++								
Basford 2000	33.3	26.3	24	56.6	61.3	24	10.3%	-23.30 [-49.99, 3.39]
Subtotal (95% CI)			**24**			**24**	**10.3%**	**-23.30 [-49.99, 3.39]**
Heterogeneity: Not applicable								
Test for overall effect: Z = 1.71 (P = 0.09)								
4.3.3 vs sham -								
Lam 2007	53.9	21.2	18	56.1	20.3	18	30.3%	-2.20 [-15.76, 11.36]
Sterigioulas_2007	25.07	10.72	25	24.25	16.04	31	59.4%	0.82 [-6.22, 7.86]
Subtotal (95% CI)			**43**			**49**	**89.7%**	**0.18 [-6.07, 6.43]**
Heterogeneity: Tau² = 0.00; Chi² = 0.15, df = 1 (P = 0.70); I² = 0%								
Test for overall effect: Z = 0.06 (P = 0.96)								
Total (95% CI)			**67**			**73**	**100.0%**	**-2.58 [-11.69, 6.52]**
Heterogeneity: Tau² = 23.39; Chi² = 2.97, df = 2 (P = 0.23); I² = 33%								
Test for overall effect: Z = 0.56 (P = 0.58)								
Test for subgroup differences: Chi² = 2.82, df = 1 (P = 0.09), I² = 64.5%