Study or Subgroup	post Mean	post SD	post Total	pre Mean	pre SD	pre Total	Weight	Mean Difference IV, Random, 95% CI
2.2.1 vs wlc +								
Subtotal (95% CI)			**0**			**0**		**Not estimable**
Heterogeneity: Not applicable								
Test for overall effect: Not applicable								
2.2.2 vs sham ++								
Basford 2000	32.1	33.6	23	27.7	25.3	23	2.6%	4.40 [-12.79, 21.59]
Subtotal (95% CI)			**23**			**23**	**2.6%**	**4.40 [-12.79, 21.59]**
Heterogeneity: Not applicable								
Test for overall effect: Z = 0.50 (P = 0.62)								
2.2.3 vs sham -								
Lam 2007	25.29	8.26	21	20.38	8.21	21	31.4%	4.91 [-0.07, 9.89]
Pettrone 2005	39.51	4.54	53	32.21	11.93	53	66.0%	7.30 [3.86, 10.74]
Subtotal (95% CI)			**74**			**74**	**97.4%**	**6.53 [3.70, 9.36]**
Heterogeneity: Tau² = 0.00; Chi² = 0.60, df = 1 (P = 0.44); I² = 0%								
Test for overall effect: Z = 4.52 (P < 0.00001)								
Total (95% CI)			**97**			**97**	**100.0%**	**6.47 [3.68, 9.26]**
Heterogeneity: Tau² = 0.00; Chi² = 0.66, df = 2 (P = 0.72); I² = 0%								
Test for overall effect: Z = 4.55 (P < 0.00001)								
Test for subgroup differences: Chi² = 0.06, df = 1 (P = 0.81), I² = 0%