Study or Subgroup	post Mean	post SD	post Total	pre Mean	pre SD	pre Total	Weight	Mean Difference IV, Random, 95% CI
2.3.1 vs wlc +								
Subtotal (95% CI)			**0**			**0**		**Not estimable**
Heterogeneity: Not applicable								
Test for overall effect: Not applicable								
2.3.2 vs sham ++								
Basford 2000	34.5	37	24	32.4	32.6	24	4.3%	2.10 [-17.63, 21.83]
Subtotal (95% CI)			**24**			**24**	**4.3%**	**2.10 [-17.63, 21.83]**
Heterogeneity: Not applicable								
Test for overall effect: Z = 0.21 (P = 0.83)								
2.3.3 vs sham -								
Lam 2007	19.56	9.75	18	18.28	9.41	18	42.2%	1.28 [-4.98, 7.54]
Pettrone 2005	36.97	14.74	54	32.89	14.74	54	53.5%	4.08 [-1.48, 9.64]
Subtotal (95% CI)			**72**			**72**	**95.7%**	**2.85 [-1.31, 7.00]**
Heterogeneity: Tau² = 0.00; Chi² = 0.43, df = 1 (P = 0.51); I² = 0%								
Test for overall effect: Z = 1.34 (P = 0.18)								
Total (95% CI)			**96**			**96**	**100.0%**	**2.81 [-1.25, 6.88]**
Heterogeneity: Tau² = 0.00; Chi² = 0.43, df = 2 (P = 0.80); I² = 0%								
Test for overall effect: Z = 1.36 (P = 0.18)								
Test for subgroup differences: Chi² = 0.01, df = 1 (P = 0.94), I² = 0%