

Supplementary Table S1. Reliability tests between “a reference” and the examiners.

Reliability (intraclass correlation coefficients (ICCs)) for opening, lateral, and protrusive movements (mm). Test 1 (2015 Jan) shows ICC values between “a reference” and the examiner who examined the first participants included in the study. Test 2 (2015 Sep), Test 3 (2017 Feb), and Test 4 (2017 Nov) are all based on ICC values between “a reference” and other examiners. The ICC values reported are average measurements. For ICC interpretation, see <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4913118/>.

	Test 1	Test 2	Test 3	Test 4
Opening movements				
Pain-free opening	0.85 (0.39,0.96)	0.96 (0.98, 0.97)	0.87 (0.84, 0.90)	0.67 (-2.14, 0.96)
Max. unassisted opening	0.93 (0.73, 0.98)	0.99 (0.98, 0.99)	0.92 (0.91, 0.94)	0.97 (0.68, 0.99)
Max. assisted opening	0.97 (0.87, 0.99)	0.99 (0.98, 0.99)	0.92 (0.91, 0.94)	0.88 (-0.16, 0.99)
Lateral at both sites and protrusive movements				
Right lateral	0.78 (-0.09, 0.96)	0.69 (0.43, 0.85)	0.89 (0.85, 0.92)	0.95 (0.53, 0.99)
Left lateral	0.83 (0.31, 0.96)	0.46 (0.34, 0.55)	0.91 (0.89, 0.93)	0.95 (0.52, 0.99)
Protrusion	0.85 (0.38, 0.96)	0.87 (0.84,0.90)	0.88 (0.83, 0.93)	0.96 (0.62, 0.99)