

Supplemental Table 1. Management of medical treatment and oral care of patients on biologics or immunosuppressants.

	All practitioners	Specialized dentist activity *	General dentist activity	<i>p-value</i>	All practitioners	Specialized dentist activity*	General dentist activity	<i>p-value</i>
	In patients under biologics				In patients under immunosuppressants			
	N = 107 (%)	N = 49 (%)	N = 58 (%)		N = 107 (%)	N = 49 (%)	N = 58 (%)	
Nature of the expected risks								
Infectious risk	105 (98.13)	49 (100.00)	56 (96.55)	0.18	106 (99.07)	49 (100.00)	57 (98.28)	0.35
Delayed healing	98 (91.59)	46 (93.88)	52 (89.66)	0.43	90 (84.11)	40 (81.63)	50 (86.21)	0.51
Hemorrhagic risk	20 (18.69)	10 (20.41)	10 (17.24)	0.67	13 (12.15)	5 (10.20)	8 (13.79)	0.57
	N = 104 (%)^a	N = 47 (%)^a	N = 57 (%)^a		N = 103 (%)^b	N = 47 (%)^b	N = 56 (%)^b	
Blood tests prescribed preoperatively								
No	80 (76.92)	32 (68.09)	48 (84.21)		74 (71.84)	31 (65.96)	43 (76.79)	
Yes	24 (23.08)	15 (31.91)	9 (15.79)	0.05	29 (28.16)	16 (34.04)	13 (23.21)	0.22
Blood cell count	23 (95.83)	15 (100.00)	8 (88.89)	0.18	28 (96.55)	16 (100.00)	12 (92.31)	0.25
Platelet cell count	17 (70.83)	11 (73.33)	6 (66.67)	0.72	14 (48.28)	8 (50.00)	6 (46.15)	0.83
Preoperative management of the medical treatment								
No modification	65 (62.50)	29 (61.70)	36 (63.16)	0.87	51 (49.51)	22 (46.81)	29 (51.79)	0.61
Discontinuation ≤ 1 month	8 (7.69)	4 (8.51)	4 (7.02)	0.77	5 (4.85)	3 (6.38)	2 (3.57)	0.50
Discontinuation > 1 month	4 (3.85)	2 (4.26)	2 (3.51)	0.84	1 (0.97)	1 (2.13)	0 (0.00)	0.27
Duration of discontinuation depending on the type of drug	3 (2.88)	2 (4.26)	1 (1.75)	0.55	1 (0.97)	1 (2.13)	0 (0.00)	0.27
Decision after consultation with the prescribing physicians	84 (80.77)	41 (87.23)	43 (75.44)	0.12	79 (76.70)	38 (80.85)	41 (73.21)	0.36
Pre, per, and post- operative dental-care protocol								
Antiseptic mouthwash								
Preoperative	74 (71.15)	33 (70.21)	41 (71.93)	0.84	69 (66.99)	31 (65.96)	38 (67.86)	0.83
Perioperative	52 (50.00)	25 (53.19)	27 (47.37)	0.55	55 (53.40)	27 (57.45)	28 (50.00)	0.45
Postoperative	79 (75.96)	37 (78.72)	42 (73.68)	0.54	74 (71.84)	32 (68.09)	42 (75.00)	0.43
Antibiotics								
Preoperative (single dose, flash)	49 (47.12)	20 (42.55)	29 (50.88)	0.39	45 (43.69)	20 (42.55)	25 (44.64)	0.83
Preoperative and postoperative (antibiotic coverage until mucosal healing)	81 (77.88)	39 (82.98)	42 (73.68)	0.25	83 (80.58)	38 (80.85)	45 (80.36)	0.95

* Specialized dentist activity: oral surgery, periodontology, endodontics, pediatric dentistry

^a Only practitioners who responded that they are concerned about the care of patients taking biologics (n = 104) were included in this analysis.

^b Only practitioners who responded that they were concerned about the care of patients taking immunosuppressants (n = 103) were included in this analysis.